




Product Sheet

Desulfovibrio desulfuricans *subsp. desulfuricans* (ATCC® 29577™)

Please read this **FIRST**



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Desulfovibrio desulfuricans subsp. desulfuricans* (ATCC® 29577™)

Description

Designation: NCIB 8307 [Essex 6]

Deposited Name: *Desulfovibrio desulfuricans subsp. desulfuricans* (Beijerinck) Kluyver and van Niel

Propagation

Medium

ATCC® Medium 1249: Modified Baar's medium for sulfate reducers

Growth Conditions

Temperature: 30.0°C

Atmosphere: Anaerobic

Propagation Procedure

1. Open vial according to enclosed instructions.
2. Perform all steps under anaerobic conditions (*see below*).
3. Aseptically transfer 0.5 ml of ATCC Medium #1249 to the vial and rehydrate the freeze-dried pellet, immediately place the rehydrated pellet under a stream of oxygen-free sterile gas. Then transfer the suspension back into the tube of broth. Inoculate a plate of non-selective medium with 0.1 of the culture. Inoculate a non-selective tube of broth.
4. Seal the test tube with a rubber stopper and incubate anaerobically at 30°C. Incubate the plate(s) and aerobic broth at 30°C as a purity check.
5. After two or three days, growth should be evident by turbidity through out the broth. Once growth has been established, the culture should be transferred to fresh broth every 24 to 48 hours.
6. This culture is very sensitive to oxygen; therefore steps should be taken to avoid exposure to oxygen. When the culture exhibits good growth it will remain viable for up to 1 week if stored at 4°C under anaerobic conditions.

ANAEROBIC CONDITIONS:

- Tubes of media are placed under a gassing cannula system hooked to a source of oxygen free gas.
- All transfers are performed while the test tubes are on the cannula system with a gentle stream of oxygen-free gas flowing through the system.
- As the test tubes are removed from the cannula system each is sealed with butyl rubber stopper thus maintaining the anaerobic headspace.

Notes

Growth is detected within 24 to 48 hours by turbidity throughout the broth.

The cells are typically comma-shaped rods that are motile.

Once growth has been established, the culture should be transferred every 24 to 48 hours when maintained at 30°C. The culture can be maintained at 4°C for up to 1 week.

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor




Product Sheet

Desulfovibrio desulfuricans
subsp. desulfuricans

(ATCC® 29577™)

Please read this **FIRST**



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Desulfovibrio desulfuricans subsp. desulfuricans* (ATCC® 29577™)

formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2014. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [12/02]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor