



Product Sheet

Leishmania donovani (ATCC® 30498™)

Please read this **FIRST**



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Leishmania donovani* (ATCC® 30498™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Strain Designation: 1S
Deposited Name: *Leishmania donovani* (Laveran and Mesnil) Ross
Depositor: R Herman
Isolation:
Bol West Village
Isolation date: 1962

Propagation

Growth Conditions
Culture System: *in vivo*, golden hamster (*Mesocricetus auratus*)
Temperature: 25°C

Protocols

Frozen ampules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampoules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

1. To thaw a frozen ampule, place it in a 35°C water bath such that the lip of the ampule remains above the water line. Thawing time is approximately 2 to 3 minutes. Do not agitate the ampule. Do not leave ampule in water bath after it is thawed.
2. Immediately after thawing, aseptically remove the entire contents of the ampule with a syringe and inject intraperitoneally into an uninfected Golden hamster at least 8 weeks old. Follow the protocol for maintenance *in vivo*. The course of infection may be longer or shorter than usual depending on percent recovery of the parasite from the frozen state.

Culture Maintenance

1. Inoculate entire infected blood suspension intraperitoneally into the hamster using a 1.0 mL syringe equipped with a 27 gauge 1/2 inch needle. Serial passage after 30-45 days.
2. To infect additional hamsters, harvest the parasites from spleen tissue homogenized in a balanced salt solution (i.e., Tyrodes' Salt solution or similar), approximately 5.0 mL salt solution per spleen. Inoculate 0.3-0.5 mL of spleen homogenate per hamster. Dosage should be a minimum of 2 x 10⁶ amastigotes per hamster. The course of infection will vary with the size of the inoculum.
3. Parasites for inoculation should be collected within 30-45 days of infection. Host hamsters should be at least 8 weeks old.

Cryopreservation

Tyrode's Salt Solution

NaCl 8.00 g
KCl 0.20 g
CaCl₂ 0.20 g
MgCl₂*H₂O 0.05 g
NaH₂PO₄*H₂O 1.00 g
NaHCO₃*H₂O 1.00 g
Glucose 1.00 g
Glass distilled H₂O to 1.00 L

Add ingredients in the sequence listed. Filter-sterilize.

1. Harvest the parasites from spleen tissue homogenized in a balanced salt solution (i.e., Tyrodes' Salt soln. or similar), approximately 5.0 mL solution per spleen.
2. Transfer the cell homogenate to a 15 mL plastic centrifuge tube and spin at approximately 1300 x g for 10 min.
3. Pool the cell pellets and adjust the concentration to 2.0 - 4.0 x 10⁷ cells/mL with a fresh solution of Tyrode's Salt Solution. *If the concentration is too low centrifuge at 1300 x g for 10 min and resuspend in the volume of Tyrode's Salt Solution required to yield the desired concentration.
4. Mix the cell preparation and 10% (v/v) DMSO in equal portions. The final concentration will be 1.0 - 2.0 x 10⁷ cells/mL and 5% DMSO. The time from the mixing of the cell preparation and cryoprotective



Product Sheet

Leishmania donovani (ATCC® 30498™)

Please read this FIRST



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Leishmania donovani* (ATCC® 30498™)

- solution to the start of the freezing process should be no less than 15 min. and no more than 30 min.
- Dispense in 0.5 mL aliquots to 1.0-2.0 mL sterile plastic screw-capped cryovials (special plastic vials for cryopreservation).
 - Place the vials in a controlled rate freezing unit. From room temperature cool at -1°C/min to -40°C. If the freezing unit can compensate for the heat of fusion, maintain rate at -1°C/min through the heat of fusion. At -40°C plunge into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately -1°C/min.)
 - Store in either the vapor or liquid phase of a nitrogen refrigerator.
 - To thaw a frozen ampule, place it in a 35°C water bath such that the lip of the ampule remains above the water line. Thawing time is approximately 2 to 3 minutes. Do not agitate the ampule. Do not leave ampule in water bath after thawed.
 - Immediately after thawing, aseptically remove the contents of the ampule with a syringe and inoculate an uninfected Golden hamster at least 8 weeks old. Follow the protocol for maintenance *in vivo*.



References

References and other information relating to this product are available online at www.atcc.org.



Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/06]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor