ATCC

Product Sheet

Campylobacter jejuni subsp. jejuni (ATCC® 43465™)

Please read this FIRST

Biosafety Level
2

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Campylobacter jejuni subsp. jejuni* (ATCC® 43465™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Designation: MK307
Deposited Name: *Campylobacter jejuni* subsp. *jejuni* (Jones et al.) Veron and Chatelain
Antigenic Properties: Serotype O:50

Propagation

Medium
ATCC® Medium 1116: Brucella broth with 0.16% agar

Growth Conditions
Temperature: 37.0°C
Atmosphere: Microaerophilic

Propagation Procedure
1. Open vial according to enclosed instructions.
2. Using a single tube of #1115, #177, or #18 broth (5 to
1. Open vial according to enclosed instructions.
2. Using a single tube of #1115, #177, or #18 broth (5 to 6 ml), withdraw approximately 0.5 to 1.0 ml with a Pasteur or 1.0 ml pipette. Rehydrate the entire pellet.
3. Aseptically transfer this aliquot back into the broth tube. Mix well.
4. Use several drops of the suspension to inoculate a #260 slant, and/or plate.
5. Or, to obtain a biphasic culture, add 0.5 ml of the suspension to a #260 agar slant (*see notes*).
6. Incubate tubes and plate at 37°C, under microaerophilic conditions, for 24 to 48 hours. Use an anaerobe jar with an active catalyst and a microaerophilic gas generator pack, or other acceptable method. Incubate slant with cap loose.

Notes

This is an organism that requires moist conditions for best growth. A biphasic culture gives the most rapid growth. Growth at the broth/agar interface of the biphasic slant should occur within one to two days, but little turbidity will be seen. To observe growth, examine a wet mount of the broth under phase microscopy. The organism is a short, thin motile rod. Motility is usually observed only in young cultures.

Growth on agar takes longer than with the biphasic culture. Once good growth is present, these organisms tend to lose viability, especially if exposed to air for lengthy periods.

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

This is an organism that requires moist conditions for best growth. A biphasic culture gives the most rapid growth. Growth at the broth/agar interface of the biphasic slant should occur within one to two days, but little turbidity will be seen. To observe growth, examine a wet mount of the broth under phase microscopy. The organism is a short, thin motile rod. Motility is usually observed only in young cultures.

Growth on agar takes longer than with the biphasic culture. Once good growth is present, these organisms tend to lose viability, especially if exposed to air for lengthy periods.

Additional information on this culture is available on the ATCC web site at www.atcc.org.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Product Sheet

Campylobacter jejuni subsp. jejuni (ATCC® 43465™)

Please read this FIRST

Biosafety Level
2

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Campylobacter jejuni subsp. jejuni* (ATCC® 43465™)

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/05]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor