ATCC

Product Sheet

# Gr431 (ATCC® HB-8575™)

## Please read this FIRST

Biosafety Level
1

## Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

## Complete Growth Medium

RPMI 1640 medium with 0.1 mM hypoxanthine and 0.016 mM thymidine, 90%; fetal bovine serum, 10%
Culture Medium: RPMI medium 1640 with 0.1 mM hypoxanthine, 0.016 mM thymidine,
90%; fetal bovine serum, 10%.

## Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: Gr431 (ATCC® HB-8575™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

*Or contact your local distributor*

## Description

**Organism:** human (B cell); mouse (myeloma)
**Isotype:** IgM
**Disease:** Cancer
**Cell Type:** hybridoma: B lymphocyte
**Morphology:** lymphoblast
**Growth Properties:** suspension

## Batch-Specific Information

Refer to the Certificate of Analysis for batch-specific test results.

## SAFETY PRECAUTION

ATCC highly recommends that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

## Unpacking & Storage Instructions

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

## Handling Procedure for Frozen Cells

Part A. FROZEN CELLS
Vol./Ampule: 1.0 mL.
Recommended Handling Upon Receipt: Initiate culture as soon as possible upon receipt. Thaw by rapid agitation in 37°C water bath. See instructions on back.
Dilute ampule contents with culture medium (see batch data above). Add fresh medium (depending on cell density) every 2-3 days.

## Handling Procedure for Flask Cultures

Part B. FLASK CULTURES
Recommended Handling Upon Receipt:
Suspension Cultures: The culture flasks have been completely filled with medium for shipment. Remove the entire contents of the flask and centrifuge at 300 x g for 15 minutes. Resuspend the cell pellet as suggested under subculture procedure described above.

## Subculturing Procedure

**Medium Renewal:** Every 2 to 3 days
Cultures can be maintained by addition or replacement of fresh medium. Start cultures at 2 X 10 exp5 cells/ml and maintain between 1 X 10 exp5 and 1 X 10 exp6 cells/ml.

## Comments

Lymphocytes were obtained from the regional lymph nodes of a patient who had breast cancer.
Lymphocytes were fused with NS-1 myeloma cells.

## References

References and other information relating to this product are available online at www.atcc.org.

## Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S.

Product Sheet

# Gr431 (ATCC® HB-8575™)

## Please read this FIRST

Biosafety Level
1

## Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

## Complete Growth Medium

RPMI 1640 medium with 0.1 mM hypoxanthine and 0.016 mM thymidine, 90%; fetal bovine serum, 10%
Culture Medium: RPMI medium 1640 with 0.1 mM hypoxanthine, 0.016 mM thymidine,
90%; fetal bovine serum, 10%.

## Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: Gr431 (ATCC® HB-8575™)

Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

## ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

## Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.
While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.
This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.
Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.
© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/06]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

*Or contact your local distributor*