

Antibiotic Susceptibility of *Escherichia coli* ATCC® BAA-3251™

Antimicrobial	MIC ^a	Interpretation ^b	Antimicrobial	MIC ^a	Interpretation ^b
Amikacin	16	S	Cefuroxime	≥64	R
Amoxicillin / Clavulanic Acid	16	I	Cefuroxime Axetil	≥64	R
Ampicillin	≥32	R	Ciprofloxacin	≥4	R
Ampicillin / Sulbactam	≥32	R	Ertapenem	≤0.5	S
Aztreonam	16	R	Gentamicin	≥16	R
Cefalotin	≥64	R	Imipenem	≤0.25	S
Cefazolin	≥64	R	Levofloxacin	≥8	R
Cefepime	2	R	Meropenem	≤0.25	S
Cefotaxime	≥64	R	Moxifloxacin	≥8	R
Cefotetan	≤4	S	Nalidixic Acid	≥32	R
Cefoxitin	≤4	S	Nitrofurantoin	≤16	S
Cefpodoxime	≥8	R	Piperacillin / Tazobactam	8	S
Ceftazidime	16	R	Tetracycline	≥16	R
Ceftazidime / Avibactam	≤0.12	S	Tigecycline	≤0.5	S
Ceftolozane / Tazobactam	≤0.25	S	Tobramycin	≥16	R
Ceftriaxone	≥64	R	Trimethoprim / Sulfamethoxazole	≥320	R

^a Antibiotic susceptibility was obtained using a Vitek2 AST-GN69 and AST-XN09 cards.

^b Parameter set: MIC Interpretation Guideline: CLSI M100-S27 (2017)
Therapeutic Interpretation Guideline: Natural Resistance