

Antibiotic Susceptibility of *Escherichia coli* ATCC® BAA-3260™

Antimicrobial	MIC ^a	Interpretation ^b	Antimicrobial	MIC	Interpretation
Amikacin	16	S	Cefuroxime	≥64	R
Amoxicillin/Clavulanic Acid	≥32	R	Cefuroxime Axetil	≥64	R
Ampicillin	≥32	R	Ciprofloxacin	≥4	R
Ampicillin/Sulbactam	≥32	R	Ertapenem	≤0.5	S
Aztreonam	≥64	R	Gentamicin	≥16	R
Cefalotin	≥64	R	Imipenem	≤0.25	S
Cefazolin	≥64	R	Levofloxacin	≥8	R
Cefepime	≥64	R	Meropenem	≤0.25	S
Cefotaxime	≥64	R	Moxifloxacin	≥8	R
Cefotetan	≥64	R	Nalidixic Acid	≥32	R
Cefoxitin	≥64	R	Nitrofurantoin	128	R
Cefpodoxime	≥8	R	Piperacillin/Tazobactam	≥128	R
Ceftazidime	≥64	R	Tetracycline	≥16	R
Ceftazidime/Avibactam	2	S	Tigecycline	≤0.5	S
Ceftolozane/Tazobactam	≥32	R	Tobramycin	≥16	R
Ceftriaxone	≥64	R	Trimethoprim/Sulfamethoxazole	≥320	R

^a Antibiotic susceptibility was obtained using a Vitek2 AST-GN69 and AST-XN09 cards.

^b Parameter set: MIC Interpretation Guideline: CLSI M100-S27 (2017)
Therapeutic Interpretation Guideline: Natural Resistance