

Antibiotic Susceptibility of *Escherichia coli* ATCC® BAA-3287™

Antimicrobial	MIC ^a	Interpretation ^b	Antimicrobial	MIC ^a	Interpretation ^b
Amikacin	16	S	Cefuroxime	≥64	R
Amoxicillin / Clavulanic Acid	16	I	Cefuroxime Axetil	≥64	R
Ampicillin	≥32	R	Ciprofloxacin	≥4	R
Ampicillin/Sulbactam	≥32	R	Ertapenem	≥8	R
Aztreonam	≥64	R	Gentamicin	≥16	R
Cefalotin	≥64	R	Imipenem	0.5	S
Cefazolin	≥64	R	Levofloxacin	≥8	R
Cefepime	≥64	R	Meropenem	≥16	R
Cefotaxime	≥64	R	Moxifloxacin	≥8	R
Cefotetan	≤4	S	Nalidixic Acid	≥32	R
Cefoxitin	8	S	Nitrofurantoin	32	S
Cefpodoxime	≥8	R	Piperacillin / Tazobactam	≥128	R
Ceftazidime	≥64	R	Tetracycline	≥16	R
Ceftazidime/Avibactam	1	S	Tigecycline	≤0.5	S
Ceftolozane/ Tazobactam	≥32	R	Tobramycin	≥16	R
Ceftriaxone	≥64	R	Trimethoprim / Sulfamethoxazole	≥320	R

^a Antibiotic susceptibility was obtained using a Vitek2 AST-GN69 and AST-XN09 cards.

^b Parameter set: MIC Interpretation Guideline: CLSI M100-S27 (2017)
Therapeutic Interpretation Guideline: Natural Resistance