

Antibiotic Susceptibility of *Escherichia coli* ATCC® BAA-3337™

Antimicrobial	MIC ^a	Interpretation ^b	Antimicrobial	MIC	Interpretation
Amikacin	8	S	Cefuroxime	≥64	R
Amoxicillin/Clavulanic Acid	16	I	Cefuroxime Axetil	≥64	R
Ampicillin	≥32	R	Ciprofloxacin	≤0.25	S
Ampicillin/Sulbactam	≥32	R	Ertapenem	≤0.5	S
Aztreonam	≤1	R	Gentamicin	≤1	S
Cefalotin	≥64	R	Imipenem	≤0.25	S
Cefazolin	16	R	Levofloxacin	≤0.12	S
Cefepime	≤1	R	Meropenem	≤0.25	S
Cefotaxime	4	R	Moxifloxacin	≤0.25	S
Cefotetan	≤4	S	Nalidixic Acid	≤2	S
Cefoxitin	≤4	S	Nitrofurantoin	≤16	S
Cefpodoxime	≥8	R	Piperacillin/Tazobactam	≤4	S
Ceftazidime	16	R	Tetracycline	≤1	S
Ceftazidime/Avibactam	≤0.12	S	Tigecycline	≤0.5	S
Ceftolozane/Tazobactam	4	I	Tobramycin	≥16	R
Ceftriaxone	8	R	Trimethoprim/Sulfamethoxazole	≥320	R

^a Antibiotic susceptibility was obtained using a Vitek2 AST-GN69 and AST-XN09 cards.

^b Parameter set: MIC Interpretation Guideline: CLSI M100-S27 (2017)
Therapeutic Interpretation Guideline: Natural Resistance