ATCC medium: 983 E medium for anaerobes with 0.2% rutin

ATCC Medium 982 (see below) with 1% phloroglucinol replaced by 0.2% rutin.

ATCC Medium 982:

Rumen Fluid (see below)	.30.0	ml
(NH ₄) ₂ SO ₄	0.5	g
Resazurin Solution (see below)	0.4	ml
Salt Solution (see below)	.50.0	ml
L-Cysteine . HCl	0.05	5 g
Sterile 1% phloroglucinol solution	.30.0	ml

Mix all the ingredients in an Erlenmeyer flask. Flask should have a small head space to minimize air volume that must be purged during cooling. Fit a removable chimney to the boiling flask to prevent medium from boiling over. Boil (10-20 min) until medium changes from pink to yellow. Replace chimney with stopper fitted with glass tubing which bubbles O_2 -free gas, composed of N_2 and H_2 through the medium to maintain in reduced state (97% N_2 , 3% H_2 is employed at the ATCC) and place in an ice bath medium into tubes under O_2 -free N_2 and N_2 . Stopper with butyl rubber stoppers. Place rack of tubes in press so that stoppers do not dislodge, and autoclave at 121C for 12-15 minutes on fast exhaust.

Rumen Fluid:

Filter rumen contents, obtained from a cow fed an alfalfa-hay concentrated ration, through two layers of cheesecloth to remove large particles. Store bottles in the freezer. Much of the particulate matter settles. Use only the supernatant.

Resazurin Solution:

Resazurin.			 •	 •		•	•	.25.0	mg
Distilled	wat	er.						100.0	ml

Salts Solution:

$CaCl_2$	(anhydrous)	g
$MgSO_4$	0.2	g
K ₂ HPO ₄	1.0	g
KH ₂ PO ₄	1.0	g
NaHCO3		g
NaCl		g

Dissolve $CaCl_2$ and $MgSO_4$ in 300 ml of distilled water. Add 500 ml water and add the remaining salts while swirling slowly. Add 200 ml of distilled water, mix, and store at 4C.

Phloroglucinol:

Filter-sterilize freshly prepared phloroglucinol solution. Wrap glassware in aluminum foil to exclude light.