



Escherichia coli bacteriophage T2

11303-B2™

Description

Escherichia coli bacteriophage T2 is a virus that infects and replicates in *Escherichia coli* strain B (ATCC 11303). This product is provided in an economical preceptrol format.

Strain designation: T2

Deposited As: T2

Storage Conditions

Product format: Frozen

Storage conditions: -80°C or colder

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submerged in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Host: *Escherichia coli* B [CIP 103914, NCIB 11595, NRC 745] (ATCC 11303)

Medium:

ATCC Medium 129: Nutrient agar/stabs/broth with 0.5% NaCl

Temperature: 37°C

Atmosphere: Aerobic

Handling Procedures

1. Carefully follow the procedures given below for phage propagation.
2. *Escherichia coli* strain B (ATCC 11303) is the recommended host.

GENERAL PROCEDURES FOR THE PROPAGATION OF BACTERIOPHAGE

To recover phage from freeze-dried or frozen vial:

Day 1

1. Prepare an actively growing culture of the recommended host strain before opening the phage specimen. The host should be 18-24 hours old.
 - a. In some instances, the host growth requirements will differ from that of the phage. Because the phage culture will only be as robust as the host culture, it is best to choose conditions according to the preference of the host bacterium.
 - b. Typically, the host will grow in the same medium as the phage, but there may be some exceptions.
2. Subculture the host on #129 agar and incubate at 37°C overnight.

Day Two

3. Based off experimental design determines the total volume needed before opening of phage.
4. Pick one colony from step b. and homogenize in 5 mL of #129 broth.
5. Incubate at 37°C in a shaking incubator (160-180 rpm) until it reaches OD₆₀₀ of 0.1 – 0.3. Use a spectrophotometer to take measurements following the manufacturer's instructions.
 - a. The culture should be checked at 1 hour and every 15 minutes thereafter until the correct measurement is achieved.
6. Thaw or rehydrate the bacteriophage seed vial. Use 0.5 mL of #129 broth to rehydrate if necessary.
7. Infect a 5 mL culture with 100 µL of the bacteriophage. Store any remaining phage at 4°C.
 - a. Perform a purity check by inoculating a blood plate and incubating at 37°C. If desired, inoculate a maintenance medium plate to check for host growth.
 - b. Inoculate a fresh subculture of the host and incubate statically at 37°C overnight.

Day Three

8. Centrifuge phage culture at 4000 g for 10 minutes. (This step is optional depending on the density of the culture it may make filtering easier). To

convert g to rpm, consult the user manual for each centrifuge. There may be a conversion tool online. 4000 g usually converts to an rpm between 6000-6500 for benchtop models.

9. Filter the lysate (supernatant) with a 0.2 to 0.45 μm PES sterile filter and store the filtrate at 4°C.
10. Prior to performing the spot titer, subculture host to a #129 broth and incubates overnight at 37°C (in case another round of amplification will be necessary the following day).
11. Melt 0.5% agar completely. The melted soft agar can be stored at 55°C for up to a week before use. Then move to the spot titer steps below.

Performing a spot titer

NOTE: Spotting the phage on plates makes visualizing the lysis easier. If phage is added directly to soft-agar before pouring plates, hazy or tiny plaques may be difficult to see. Resistant host bacteria may also mask plaque formation.

12. To perform a spot titer, warm one or two plates at 37°C. The soft agar should be brought to 43°C to 45°C until ready to pour. It may be advisable to use a water bath.
13. Overlay the surface with 2.5 mL of melted 0.5% agar containing 100 μL of the overnight host culture. Allow overlay to harden. This usually won't take longer than 15 to 30 minutes.
14. The phage lysate can be serial diluted in a 96 well plate in quadruplicate (if desired). Aliquot 90 μL of #129 broth medium into each well.
15. Add 10 μL of phage lysate to each well and mix. Pass 10 μL to each of the next set of wells and mix. Continuing to the desired number of passages.
16. Spot 2 μL of each dilution on the plate from step o. Up to 8 dilutions can fit on a 90 mm petri dish. Incubate plates at 37°C, inverted is not recommended as it can affect the spots that are spotted. Note: If over-incubated, plaques can continue to enlarge. Given sufficient numbers of plaques, this may result in confluent lysis.

Day Four

This could be a stopping point depending on the total volume achieved from the first round of amplification and the total volume needed for experimentation. Count the plaques in each spot of the dilution with the best growth. Calculate the average count. Use the formula below to determine the pfu/mL.

$$\text{a. pfu/mL} = \text{average plaque count} / [(\text{dilution factor}) (2 \times 10^{-3} \text{mL})]$$

To propagate phage:

17. Determine the total volume needed and place this amount of broth in a flask. Add a small amount of overnight host culture to the flask and incubate at 37°C while shaking until the growth reaches OD₆₀₀ of 0.1 – 0.3.
18. Infect with the calculated volume of phage lysate using the following formula. Volume of phage to add (ml) = $(8 \times 10^8 \times \text{total culture volume in ml} \times \text{OD}_{600} \times \text{MOI}) / \text{phage titer (PFU/ml)}$. Shake at 160-180 rpm at 37°C overnight.
19. Centrifuge phage culture at 4000 g for 10 minutes. Filter the lysate with a 0.2 µm or 0.45 µm PES sterile filter. The filtration can be stored at 4°C.
20. Lysates should remain viable under refrigeration for long periods. They may also be frozen with or without cryoprotectant. If available, liquid nitrogen storage is the best method for long term storage. Most phages can also be freeze-dried. ATCC® uses double strength skim milk mixed half and half with the filtrate.
21. Lysates should remain viable under refrigeration for long periods. Storage at 20°C may cause inactivation of the phage. They may be frozen with cryoprotectant. If available, liquid nitrogen storage is the best method for long term storage. Most phages can also be freeze-dried. ATCC® uses double strength skim milk mixed half and half with the filtrate.

Notes

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Escherichia coli* bacteriophage T2 (ATCC 11303-B2)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2026-06-25

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

***Escherichia coli* bacteriophage T2**

11303-B2

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
