



Desulfovibrio salexigens Postgate and Campbell

14944™

Description

Strain designation: NCIB 8365 [Louisiana 43.11]

Deposited As: *Desulfovibrio salexigens* Postgate and Campbell

Type strain: No

Storage Conditions

Product format: Freeze-dried

Storage conditions: 2°C to 8°C

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submerged in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Medium:

ATCC Medium 1250: Modified Barr's Medium for sulfate reducers with 2.5% NaCl

Temperature: 37°C

Atmosphere: 80% N₂, 20% CO₂

Incubation: With shaking

Handling Procedures

1. Open vial according to enclosed instructions.
2. Perform all steps under anaerobic conditions. (*see below*). Exchange the gas in the head space for a fresh anaerobic gas, either 80% N₂ 20% CO₂ or 100% N₂. To insure that the media is anaerobic add 0.1 mL 1.5% sodium sulfide (stock concentration) for each 5 to 10 mL of medium.

14944

3. Using an anaerobic 1 mL syringe (see below) aseptically transfer 0.5 mL of ATCC Medium #1250 to the vial and rehydrate the entire freeze-dried pellet. Transfer the entire suspension back into the primary tube. The primary tube should not contain more than 8 to 10 mL of #1250 broth. Secondary tube(s) can be inoculated with 0.5 mL of the primary broth. Inoculate a plate of non-selective medium with 0.1 of the culture.
4. Incubate the tubes anaerobically at 37°C with gentle shaking. Incubate the plate(s) aerobically as a purity check.
5. After two or three days, growth should be evident as indicated by turbidity throughout the broth. Once growth has been established, the culture should be transferred to fresh broth every 24 to 48 hours.
6. This culture is sensitive to oxygen, therefore steps should be taken to avoid exposure to oxygen. When the culture exhibits good growth it will remain viable for up to 1 week if stored at 4°C under anaerobic condition.

ANAEROBIC CONDITIONS:

- A. Balch tubes (available from Bellco Glass, Vineland, NJ; are specially designed for anaerobic work and use an aluminum crimp cap to hold a rubber stopper in place. Needles can easily be inserted through the stopper. Alternatively, serum vials may be used, or screw cap tubes with butyl rubber stoppers, in the latter case the stopper may be removed and the tube placed under a cannula system that dispenses sterile, oxygen free gas for the addition of reducing agents or inoculation.
- B. To obtain a fully reduced medium, it is necessary that the medium be anoxic and that a reducing agent be added. Common reducing agents are sodium sulfide, cysteine, dithiothreitol, titanium citrate and Co-enzyme M (see D). If component IV is added to the medium sodium sulfide, dithiothreitol and titanium citrate will cause the ferrous ammonium sulfate to precipitate even without growth.
- C. We suggest adding the reducing agent to the medium at least one hour before the medium is to be inoculated.
- D. Syringes can be made anaerobic by one of two methods.
 1. Displace the dead space in the syringe with a sterile oxygen-free gas.
 2. Displace the dead space in the syringe with a reducing agent.

Notes

Growth should be detected within 24 to 48 hours as indicated by turbidity throughout the broth. Turbidity may not be evident at 24 hours in the secondary growth tubes, but viable cells can be observed microscopically.

The cells typically appear as comma-shaped rods that are motile.

Once growth has been established the culture should be transferred every 24 hours when maintained at 37°C. The culture can be maintained at 4°C for up to 1 week.

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Desulfovibrio salexigens* Postgate and Campbell (ATCC 14944)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium

formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the

***Desulfovibrio salexigens* Postgate and Campbell**

14944

use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2026-06-28

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
