

50092<sup>™</sup>

### **Description**

Strain designation: NEPCC 242 [ATCC 50094] Deposited As: Pavlova lutheri (Droop) Green

Type strain: No

### **Storage Conditions**

Product format: Test tube

Storage conditions: See handling procedure

### Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

### BSL<sub>1</sub>

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of Biosafety in Microbiological and Biomedical Laboratories (BMBL), U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.



ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

### Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

### **Growth Conditions**

Medium:

ATCC Medium 1405: HESNW medium

Instructions for complete medium: Grown with mixed bacteria

**Temperature:** 20-25°C **Atmosphere:** Aerobic

## Handling Procedures

### **Handling of Live Culture**

This strain is routinely shipped as a growing culture in a glass 16 x 125 mm screw-capped test tube. The volume of the cell suspension is approximately 5-6 mL. When the culture arrives remove it promptly from the shipping container. **Do not store the** 

culture at refrigeration temperatures before handling. To assure viability, immediately loosen the test tube cap and incubate under light (~50 mEinsteins/m²/s irradiance) at a 15° horizontal slant at 20-25°C for at least one hour before observing the culture. There should be numerous active trophozoites in suspension. If the numbers are low the culture may have been exposed to temperature extremes in transit. Regardless of the state of the culture, seal and agitate the culture, then aseptically transfer a 0.5 mL aliquot to a 16 x 125 mm screw-capped test tube containing 5 mL of sterile ATCC medium 1405. Incubate the parent and daughter cultures at a 15° horizontal slant at 20-25°C with the caps on loosely for air exchange. Maintain under a 14 hour light (~50 mEinsteins/m²/s irradiance)/10 hour dark cycle.

#### **Culture maintenance:**

Subculture every two weeks to a fresh tube of medium in the following manner:

- Seal and agitate the parent culture to suspend cells, then aseptically transfer
  mL to a 16 x 125 mm screw-capped test tube containing 5 mL of sterile
  ATCC medium 1405.
- 2. Incubate the parent and daughter cultures at a 15° horizontal slant at 20-25°C with the caps on loosely for air exchange. Maintain under a 14 hour light (~50 mEinsteins/m²/s irradiance)/10 hour dark cycle.

**Note:** Addition of penicillin G at 50-75 U/mL and streptomycin at 50-75  $\mu$ g/mL to the culture may be necessary if the bacterial density in the culture becomes high enough to affect growth.

### Reagents for cryopreservation: <u>Cryoprotective Solution</u>

Glycerol 2.4 mL

Fresh growth medium w/o bacteria 7.6 mL

### **Cryopreservation:**

- 1. Mix the components of the cryoprotective solution in the order listed.
- 2. Harvest cells from a culture that is at or near peak density by centrifugation at  $800-1000 \times q$  for 5 min.
- 3. Adjust the concentration of cells to at least  $2 \times 10^7$ /mL in fresh medium.
- 4. Mix the cell preparation and the cryoprotective solution in equal portions.
- 5. Dispense in 0.5 mL aliquots into 1.0 2.0 mL sterile plastic screw-capped cryules (special plastic vials for cryopreservation).
- 6. Place vials in a controlled rate freezing unit. From room temperature cool at -



1°C/min to -40°C. If freezing unit can compensate for the heat of fusion, maintain rate at -1 C/min through heat of fusion. At -40°C plunge ampules into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately -1°C/min.)

- 7. Ampules are stored in either the vapor or liquid phase of a nitrogen refrigerator.
- 8. To establish a culture from the frozen state place the vial in a 35°C water bath. Immerse the vial to a level just above the surface of the frozen material. Do not agitate the vial. Immediately after thawing, do not leave in water bath, aseptically remove the contents of the ampule and inoculate into a 16 x 125 mm screw-capped test tube containing 5 mL of sterile ATCC medium 1405. Immediately seal and agitate the culture to evenly suspend cells, then aseptically transfer a 0.5 mL aliquot to a second, identical tube of medium.
- 9. Incubate the parent and daughter cultures at a 15° horizontal slant at 20-25°C with the caps on loosely for air exchange. Maintain under a 14 hour light ( $\sim$ 50 µEinsteins/m²/s irradiance)/10 hour dark cycle.
- 10. Follow the protocol for maintenance of culture.

### Notes

This xenic culture contains the original bacterial flora present when the alga was first isolated.

### Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Pavlova lutheri* (Droop) Green (ATCC 50092)

#### References

References and other information relating to this material are available at



www.atcc.org.

### Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

### **Disclaimers**

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and

assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

## Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

### Revision

This information on this document was last updated on 2021-05-19

### Contact Information

**ATCC** 

10801 University Boulevard Manassas, VA 20110-2209

**USA** 

US telephone: 800-638-6597



Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor

