



Balamuthia mandrillaris

Visvesvara et al.

50209TM

Description

This parasitic protozoan was isolated from the brain tissue of a mandrill that died of amebic meningoencephalitis in the San Diego Zoo.

Strain designation: CDC:V039

Deposited As: *Balamuthia mandrillaris* Visvesvara et al.

Type strain: Yes

Storage Conditions

Product format: Frozen

Storage conditions: -80°C or colder for 1 week, vapor phase of liquid nitrogen for long-term storage

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories* (BMBL), U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local



***Balamuthia mandrillaris* Visvesvara et al.**

50209

or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Host: VERO C1008 [Vero 76, clone E6, Vero E6] (ATCC CRL-1586); Vero (ATCC CCL-81)

Medium:

ATCC Medium 2222: Cell Cultivation Medium for Parasites

Instructions for complete medium: Eagle's Minimum Essential Medium (EMEM) (ATCC 30-2003) with 2 mM L-glutamine and Earle's BSS adjusted to contain 1.5 g/L sodium bicarbonate, 0.1 mM non-essential amino acids, and 1.0 mM sodium pyruvate; supplemented with 3% fetal bovine serum (ATCC 30-2020).

Temperature: 35-37°C

Handling Procedures



ATCC® Credible leads to Incredile®

www.atcc.org

Cell Line Maintenance:

1. To establish a cell culture from the frozen state place an ampule in a water bath set at 35°C (2-3 min). Immerse the vial just sufficiently to cover the frozen material. Do not agitate the vial.
2. Immediately after thawing, aseptically remove the contents of the ampule and inoculate into 10.0 ml of fresh ATCC 30-2003 with 10% (v/v) Heat-Inactivated Fetal Bovine Serum (HIFBS)* in a T-25 tissue culture flask.
3. Outgas the flask for 10 seconds with a 95% air, 5% CO₂ gas mixture.
4. Incubate in a 35°C-37°C CO₂ incubator with the cap screwed on tightly.
5. Change the medium 1-2 times per week.

* Fetal bovine serum is available from ATCC (ATCC 30-2020). Serum is heat-inactivated by exposure to 56°C for 30 minutes. This treatment will inactivate proteins of the complement pathway. Remove the serum from the refrigerator and aseptically distribute in 100 ml aliquots to sterile 125 ml screw-capped bottles. Immerse bottles in a 35°C water bath for 5 minutes. Do not directly transfer bottles from the refrigerator to 56°C. Transfer the bottles to a 56°C water bath and begin timing for 30 minutes. To avoid contamination, do not allow the level of the water in the bath to come in contact with the lip of the screw cap. It is best to leave one inch between the serum level in the bottle and the lip of the cap and to fill the water bath to a level just slightly above the level of the serum. To assure even heating of the serum, swirl the bottle(s) every ten minutes. Note: Some suppliers provide serum already heat-inactivated.

Transferring the Cell Line:

1. When the cell line forms a confluent layer, remove all the medium and replace it with 3 ml of Phosphate Buffered Saline (PBS) (ATCC 30-2200). Incubate T-25 flask at 35°C-37°C for 25-30 min.
2. Remove all the PBS and replace it with 2 ml of 0.25% (w/v) trypsin dissolved in Hank's Balanced Salt Solution (ATCC 30-2101).
3. Gently distribute the trypsin over the monolayer, remove the trypsin, and place the flask at 35°C-37°C for 10 min.
4. Add 2 ml of ATCC 30-2003 with 10% (v/v) HIFBS and detach any cells still adherent by alternately aspirating the medium into a pipette and discharging the contents over the monolayer.
5. Distribute the cell suspension in 0.5 ml aliquots to 4 T-25 flasks containing 10 ml fresh ATCC 30-2003 with 10% (v/v) HIFBS.

Balamuthia mandrillaris Visvesvara et al.

50209

6. Outgas the flask for 10 seconds with a 95% air, 5% CO₂ gas mixture.
7. Incubate in a 35°C-37°C CO₂ incubator with the cap screwed on tightly.

Establishing a *Balamuthia* Culture from the Frozen State:

Frozen ampules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampoules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

1. To thaw a frozen ampule, place it in a 35°C water bath such that the lip of the ampule remains above the water line. Thawing time is approximately 2 to 3 minutes. Do not agitate the ampule. Do not leave ampule in water bath after it is thawed.
2. Immediately after thawing, aseptically transfer contents to a T-25 tissue culture flask containing a fresh monolayer of cells (ATCC CRL-1586 or ATCC CCL-81) and 10 ml ATCC 30-2003 with 3% (v/v) HIFBS.
3. Outgas the flask for 10 seconds with a 95% air, 5% CO₂ gas mixture.
4. Incubate in a 35°C-37°C CO₂ incubator with the cap screwed on tightly.

Culture maintenance:

1. Allow the *Balamuthia* to completely destroy the host cell monolayer and encyst.
2. Remove the medium from a fresh confluent monolayer of cells (ATCC CRL-1586 or ATCC CCL-81) in a T-25 tissue culture flask and replace it with 10 ml of ATCC 30-2003 with 3% (v/v) HIFBS.
3. To transfer the *Balamuthia* culture, use a sterile cell scraper to detach cysts from the *Balamuthia* culture, agitate gently to evenly suspend the cysts, and aseptically transfer a 0.5 ml aliquot to the flask of cells prepared in step 2.
4. Outgas the flask for 10 seconds with a 95% air, 5% CO₂ gas mixture.
5. Incubate in a 35°C-37°C CO₂ incubator with the cap screwed on tightly.

Cryopreservation:

1. To harvest the *Balamuthia* culture, detach cysts by scraping the inside bottom surface of the flask with a sterile cell scraper.
2. Transfer the cyst suspension to 15 ml plastic centrifuge tubes. Centrifuge at approximately 800 x g for 5 min.
3. Remove all but 0.5 ml of the supernatant from each tube, resuspend the cyst pellets, and pool them to a single tube.
4. Adjust the concentration of cysts to 2.0 - 4.0 x 10⁵ cysts/ml with fresh medium or PBS.

***Balamuthia mandrillaris* Visvesvara et al.**

50209

NOTE: If the concentration of cysts is too low, centrifuge at 800 x g for 5 min and resuspend in the volume of fresh medium or PBS required to yield the desired concentration.

5. Prepare a cryoprotective solution containing 15% (v/v) DMSO and 6% (v/v) HIFBS in fresh medium or PBS.
6. Mix the cyst preparation and cryoprotective solution in equal portions. The final concentration will be $1.0 - 2.0 \times 10^5$ cysts/ml, 7.5% DMSO, and 3% HIFBS. The time from the mixing of the cyst preparation and cryoprotective solution to the start of the freezing process should be no less than 15 min. and no more than 30 min.
7. Dispense in 0.5 ml aliquots to 1.0-2.0 ml sterile plastic screw-capped cryules (special plastic vials for cryopreservation).
8. Place vials in a controlled rate freezing unit. From room temperature cool at $-1^{\circ}\text{C}/\text{min}$ to -40°C . If freezing unit can compensate for the heat of fusion, maintain rate at $-1^{\circ}\text{C}/\text{min}$ through heat of fusion. At -40°C plunge ampules into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately $-1^{\circ}\text{C}/\text{min}$.)
9. Store frozen ampules in either the vapor or liquid phase of a nitrogen refrigerator.
10. To thaw a frozen ampule, place it in a 35°C water bath such that the lip of the ampule remains above the water line. Thawing time is approximately 2 to 3 minutes. Do not agitate the ampule. Do not leave ampule in water bath after thawed.
11. Immediately after thawing, aseptically transfer contents to a T-25 tissue culture flask containing a fresh monolayer of cells (ATCC CRL-1586 or ATCC CCL-81) and 10 ml ATCC 30-2003 with 3% (v/v) HIFBS.
12. Outgas the flask for 10 seconds with a 95% air, 5% CO_2 gas mixture.
13. Incubate in a 35°C - 37°C CO_2 incubator with the cap screwed on tightly.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Balamuthia mandrillaris* Visvesvara et al. (ATCC 50209)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility

of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-08-21

Contact Information

ATCC

10801 University Boulevard



ATCC® Credible leads to Incredible®

www.atcc.org

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
