



Entamoeba histolytica Schaudinn

Product Sheet

50412TM

Description

Strain designation: SAW 891 R Clone B

Deposited As: *Entamoeba histolytica* Schaudinn

Type strain: No

Storage Conditions

Product format: Frozen

Storage conditions: -80°C or colder for 1 week, vapor phase of liquid nitrogen for long-term storage

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of the *Guidelines for Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For vials that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may cause the vials to

Entamoeba histolytica Schaudinn

50412

vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, do not allow these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Medium:

ATCC Medium 2154: LYI Entamoeba medium

Instructions for complete medium:

ATCC Medium PRA-2154

(Quality controlled freeze-dried lots of this medium are commercially available from ATCC).

Temperature: 35°C**Atmosphere:** Anaerobic**Culture system:** Axenic

Handling Procedures

Storage and Culture Initiation

Frozen ampules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen facilities are not available, frozen ampules may be stored at or below -70°C for approximately one week. **Do not in any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen ampules at this temperature will result in the death of the culture.

1. To thaw a frozen ampule, place in a 35°C water bath, until thawed (2-3 min). Immerse the ampule just so as to cover the frozen material. Do not agitate the ampule.

***Entamoeba histolytica* Schaudinn**

50412

2. Immediately after thawing, aseptically transfer contents to a glass screw-capped tube containing 2154. Screw cap on tightly and incubate on a 15° horizontal slant at 35°C.

Culture maintenance:

1. Ice culture at or near peak density for 10 min.
2. Gently invert culture 20 times.
3. Aseptically transfer a 0.1 and 0.25 ml aliquot to freshly prepared (no older than 7-10d) tubes of ATCC medium 2154.
4. Screw caps on tightly and incubate at a 15° horizontal slant at 35°C.
5. Subculture every 10-14 days.

Reagents for cryopreservation: CPMB-5 Cryoprotective Solution

DMSO	1.0 ml
2.5 M Sucrose	0.8 ml
L-Cysteine/Ascorbic Acid Solution	0.2 ml
CPMB-2 Basal Solution	6.0 ml
HIBS	2.0 ml

CPMB-2 Basal Solution

Yeast Extract 60.0 g

K₂HPO₄ 1.0 g

KH₂PO₄ 0.6 g

NaCl 2.0 g

Distilled water 1.0 L

Autoclave for 15 minutes.

L-Cysteine/Ascorbic Acid Solution

L-Cysteine-HCl 1.0 g

Ascorbic Acid 0.1 g

Distilled water 10.0 ml

Add 9.0 ml of distilled water to a 20 ml beaker and dissolve the first two components. While stirring, adjust the pH to 7.0 with 10N NaOH (approximately 0.7 ml). Adjust final volume to 10 ml with distilled water and filter sterilize. Solution is stable for 1 month. Discard any unused solution.

Cryopreservation:

1. Harvest cells from several cultures that are in the late logarithmic to early stationary phase of growth. Harvest cells into 50 ml of 0.22 µm filtered medium and centrifuge at 1000 x g for 10 min.



***Entamoeba histolytica* Schaudinn**

50412

2. Invert tubes 20 times and centrifuge at 200 x g for 5 min.
3. While cells are centrifuging, prepare the cryoprotective solution.
 - a) Place 1.0 ml of DMSO in a 16 x 125 mm screw-capped test tube and ice until solidified.
 - b) Add 0.8 ml of the 2.5 M Sucrose solution, remove from ice and invert until the DMSO is liquefied. Re
 - c) Add 0.2 ml of the L-Cysteine/Ascorbic Acid Solution to the DMSO solution and mix.
 - d) Add 6.0 ml of the CPMB-2 Basal Solution and mix. e) Add 2.0 ml HIBS and mix.
4. Resuspend the cell pellets and pool to a final volume of approximately 10 ml with the supernatant. Make determination of the cell density and adjust the concentration of the cells between 5×10^5 /ml - 1×10^6 medium. If the cell concentration is below 5×10^5 /ml, centrifuge the cell suspension and resuspend the volume that will yield the desired concentration.
5. After the cell concentration is adjusted, centrifuge as in step 2.
6. Remove as much supernatant as possible and determine the volume removed.
7. Resuspend the cell pellet with a volume of the cryoprotective solution equal to the volume of the supernatant removed. Invert the tube several times to obtain a uniform cell density.
8. Dispense 0.5 ml aliquots into 1.0 - 2.0 ml plastic sterile cryules (special plastic vials for cryopreservation).
9. Place the vials in a controlled rate freezing unit. Use the following cooling cycle: From room temperature at $10^{\circ}\text{C}/\text{min}$ to the heat of fusion; from the heat of fusion to -40°C , cool at $-1^{\circ}\text{C}/\text{min}$. At -40°C plunge into liquid nitrogen. The cooling cycle should be initiated no less than 15 and no more than 30 minutes after the addition of cell preparation.
10. Store ampules in a liquid nitrogen refrigerator until needed.
11. To establish a culture from the frozen state, place an ampule in a 35°C water bath, until thawed (2-3 minutes). Add a small amount of medium to the ampule, just sufficient to cover the frozen material. Do not agitate the ampule.
12. Transfer contents of thawed ampule to a 16 x 125 mm screw-capped borosilicate glass test tube containing ATCC medium 2154.
13. Screw cap on tightly and incubate at a 15° horizontal slant at 35°C . Observe the culture daily and transfer to a new tube when trophozoites are observed.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Entamoeba* Schaudinn (ATCC 50412)

References

Entamoeba histolytica Schaudinn

50412

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of sale provided that the customer has stored and handled the product according to the information included on the information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and conditions that have been found to be effective for the product. While other unspecified media and reagents may also produce similar results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and viability of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer in effect as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate and complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility associated with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable efforts are made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the use or misrepresentation of such materials.

***Entamoeba histolytica* Schaudinn**
50412

Please see the material transfer agreement (MTA) for further details regarding the use of this product.
www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-08-21

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
