**Product Sheet** 

## Giardia intestinalis (Lambl) Alexeieff

**50585**<sup>™</sup>

#### Description

**Strain designation:** AB **Deposited As:** *Giardia intestinalis* (Lambl) Alexeieff **Type strain:** No

## **Storage Conditions**

**Product format:** Frozen **Storage conditions:** -80°C or colder for 1 week, vapor phase of liquid nitrogen for long-term storage

## Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

## BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.



www.atcc.org

Page 1 of 7

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

## **Certificate of Analysis**

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

### **Growth Conditions**

Medium: ATCC Medium 2695: Keister's Modified TYI-S-33 ATCC Medium 2155: LYI Giardia Medium (filtered) Instructions for complete medium: Media: ATCC Medium 2695† (Quality controlled freeze-dried lots of this medium are commercially available as cat. no. PRA-2695) †previously ATCC Medium 1404

Alternate Media: ATCC Medium 2155 (Quality controlled freeze-dried lots of Temperature: 35°C Culture system: Axenic

Handling Procedures



50585

#### **Storage and Culture Initiation**

Frozen ampules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

- 1. To thaw a frozen ampule, place it in a 35°C water bath , until thawed (2-3 min). Immerse the ampule just sufficient to cover the frozen material. Do not agitate the ampule.
- Immediately after thawing, aseptically transfer contents to a screw-capped test tube containing 13 mL ATCC Medium 2695. Incubate the tube on a 15° horizontal slant at 35°C.

#### Culture maintenance:

- 1. When the culture has reached or is near peak density, place the tubes on ice for 10 minutes.
- 2. Gently invert the culture tube 10 times and aseptically transfer a 0.1-0.4 mL aliquot to a screw-capped test tube containing 13 mL ATCC Medium 2695.
- 3. Incubate the culture on a 15° horizontal slant at 35°C.
- 4. Transfer the culture every 3-4 days as described in steps 1-2. The transfer interval will depend on the size of the inoculum and the quality of the medium. This should be empirically determined by examining the culture on a daily basis until the growth cycle has stabilized. Do not allow the culture to overgrow. The culture crashes soon after reaching peak density.

#### 5. Cryopreservation:

- Harvest cells from a culture that is at or near peak density. To detach cells from the wall of the culture tubes place on ice for 10 minutes. Invert tubes several times until the majority of the cells are in suspension. Centrifuge tubes at 800 x g for 5 minutes.
- 2. Adjust the concentration of cells to 2 x  $10^7$ /mL in fresh medium.
- 3. Before centrifuging prepare a 24% (v/v) solution of sterile DMSO in fresh medium containing 8% (w/v) sucrose. The solution is prepared as follows:
  - a. Add 1.05 g sucrose to 10 mL of fresh medium and filter sterilize through a 0.2  $\mu m$  filter;
  - b. Add 2.4 mL of DMSO to an ice cold 20 x 150 mm screw-capped test tube;
  - c. Place the tube on ice and allow the DMSO to solidify (~5 min) and

then add 7.6 mL of ice cold medium prepared in step 3a. The final concentration will be 24% (v/v) DMSO and 8% (w/v) sucrose;

- d. Invert several times to dissolve the DMSO;
- e. Allow to warm to room temperature.
- 4. Mix the cell preparation and the cryoprotective agent, prepared in step 3, in equal portions. Thus, the final concentration will equal 12% (v/v) DMSO + 4% sucrose (w/v) and 10<sup>7</sup> cells/mL. The time from the mixing of the cell preparation and DMSO stock solution to the start of the freezing process should be no less than 15 min and no longer than 30 min.
- 5. Dispense in 0.5 mL aliquots into 1.0 2.0 mL sterile plastic screw-capped cryules (special plastic vials for cryopreservation).
- 6. Place the vials in a controlled rate freezing unit. From room temperature cool at -1°C/min to -40°C. If the freezing unit can compensate for the heat of fusion, maintain rate at -1°C/min through the heat of fusion. At 40°C plunge into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately -1°C/min.)
- 7. The frozen preparations should be stored in either the vapor or liquid phase of a nitrogen refrigerator. Frozen preparations stored below -130°C are stabile indefinitely. Those stored at temperatures above -130°C are progressively less stabile as the storage temperature is elevated.
- 8. To establish a culture from the frozen state place an ampule in a water bath set at 35°C. Immerse the vial just to a level just above the surface of the frozen material. Do not agitate the vial.
- 9. Immediately after thawing, do not leave in the water bath, aseptically remove the contents of the ampule and inoculate a 16 x 125 mm screw-capped test tube containing 13 mL ATCC Medium 2695.
- 10. Incubate the culture on a 15° horizontal slant at 35°C.

#### **Material Citation**

If use of this material results in a scientific publication, please cite the material in the following manner: *Giardia intestinalis* (Lambl) Alexeieff (ATCC 50585)



#### 50585

#### References

References and other information relating to this material are available at www.atcc.org.

#### Warranty

The product is provided 'AS IS' and the viability of ATCC<sup>®</sup> products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

#### Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility



www.atcc.org

of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

#### Copyright and Trademark Information

© ATCC 2023. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection.

#### Revision

This information on this document was last updated on 2024-10-24

#### **Contact Information**

ATCC 10801 University Boulevard



50585

Manassas, VA 20110-2209 USA US telephone: 800-638-6597 Worldwide telephone: +1-703-365-2700 Email: tech@atcc.org or contact your local distributor



