Product Sheet

🜵 Subulatomonas tetraspora

50623[™]

Description

Strain designation: plma-1 Deposited As: Subulatomonas tetraspora Type strain: No

Storage Conditions

Product format: Frozen **Storage conditions:** -80°C or colder for 1 week, vapor phase of liquid nitrogen for long-term storage

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always



used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Medium:

ATCC Medium 1989: Seawater Medium for Selected Anerobic Protists **Instructions for complete medium:** ATCC Medium 1989 inoculated with *Klebsiella pneumoniae* subsp. *pneumoniae* (ATCC[®] 700831) or *Enterobacter aerogenes* (ATCC[®] 13048). **Temperature:** 25°C **Atmosphere:** Microaerophilic **Incubation:** Grown with Enterobacter aerogenes ATCC 13048 or Klebsiella pneumoniae subsp. pneumoniae ATCC 700831 as a food source.

Handling Procedures

Storage and Culture Initiation:

Frozen ampules packed in dry ice should either be thawed immediately or stored in

Subulatomonas tetraspora

50623

liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampoules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

- One day before thawing the ampule, prepare bacterized ATCC medium 1989, i.e. inoculate medium with a bacteriological loop of *Klebsiella pneumoniae* subsp. *pneumoniae* (ATCC[®] 700831[™]) or *Enterobacter aerogenes* (ATCC[®] 13048[™]) from a nutrient agar slant (ATCC medium 3).
- 2. To thaw a frozen ampule, place it in a 35°C water bath such that the lip of the ampule remains above the water line. Thawing time is approximately 2 to 3 minutes. Do not agitate the ampule. Do not leave ampule in water bath after it is thawed.
- 3. Immediately after thawing, aseptically transfer contents to a 16 x 125 mm screw-capped test tube containing 13 mL of bacterized ATCC Medium 1989. Screw the cap on tightly and incubate upright at 25°C.

Culture maintenance:

- 1. Prepare bacterized ATCC medium 1989.
- 2. When the culture is at or near peak density, rub the surface of the tube with a sterile cotton swab, and agitate the swab to dislodge the adherent cells. Invert gently 10 times to distribute cells evenly.
- 3. Transfer approximately 0.25 mL to a 16 x 125 mm screw-capped test tube containing 13 mL of fresh, bacterized ATCC medium 1989.
- 4. Screw the cap on tightly and incubate upright at 25°C.

Cryopreservation:

- 1. Harvest the cells from a culture that is at or near peak density by centrifuging at 850 x g for 5 minutes.
- 2. If the cell concentration exceeds the required level do not centrifuge, but adjust the concentration to between 2×10^6 and 2×10^7 cells/mL with fresh medium. If the concentration is too low, centrifuge at 850 x g for 5 min and resuspend the pellet in the volume of fresh medium required to yield the desired concentration.
- 3. While cells are centrifuging prepare a 20% (v/v) solution of sterile DMSO as follows: Add the required volume of DMSO to a glass screw-capped test tube and place it in an ice bath. Allow the DMSO to solidify. Add the required volume of refrigerated medium. Dissolve the DMSO by inverting the tube several times. *NOTE: If the DMSO solution is not prepared on ice, an exothermic

50623

reaction will occur that may precipitate certain components of the medium.

- 4. Mix the cell preparation and the DMSO in equal portions. Thus, the final concentration will be between 10^6 and 10^7 cells/mL and 10% (v/v) DMSO. The time from the mixing of the cell preparation and DMSO stock solution to start of the freezing process should be no less than 15 min and no longer than 30 min.
- 5. Dispense in 0.5 mL aliquots into 1.0 2.0 mL sterile plastic screw-capped cryules (special plastic vials for cryopreservation).
- 6. Place vials in a controlled rate freezing unit. From room temperature cool at -1°C/min to -40°C. If freezing unit can compensate for the heat of fusion, maintain rate at -1°C/min through heat of fusion. At -40°C plunge ampules into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately -1°C/min.)
- 7. The frozen preparations are stored in either the vapor or liquid phase of a nitrogen freezer.
- 8. To establish a culture from the frozen state place an ampule in a water bath set at 35°C (2-3 min). Immerse the vial just sufficient to cover the frozen material. Do not agitate the vial.
- Immediately after thawing, aseptically remove the contents of the ampule and inoculate into 13 mL of fresh, bacterized ATCC medium 1989 in a 16 x 125 screw-capped test tube. Incubate upright at 25°C.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Subulatomonas tetraspora* (ATCC 50623)

References

References and other information relating to this material are available at www.atcc.org.



Warranty

The product is provided 'AS IS' and the viability of ATCC[®] products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity



www.atcc.org

Page 5 of 6

undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-07-18

Contact Information

ATCC 10801 University Boulevard Manassas, VA 20110-2209 USA US telephone: 800-638-6597 Worldwide telephone: +1-703-365-2700 Email: tech@atcc.org or contact your local distributor

