

51918<sup>™</sup>

### Description

**Strain designation:** N-5 [Glencoe 2]

Deposited As: Nitrobacter sp.

Type strain: No

## **Storage Conditions**

**Product format:** Frozen

Storage conditions: -80°C or colder

#### Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

#### BSL<sub>1</sub>

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories* (*BMBL*), U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is



51918

important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

## Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

#### **Growth Conditions**

**Medium:** 

ATCC Medium 480: Nitrobacter medium 203

**Temperature:** 26°C **Atmosphere:** Aerobic

## Handling Procedures

- 1. Allow vial to thaw. Transfer the entire contents into 5.0 mL of #480 broth contained in a large test tube (20 X 150 mm) or a small flask.
- 2. Incubate tube at 26°C, static, in the dark, and in a slanted position to increase surface area.
- 3. Monitor culture growth daily by assaying for the disappearance of NO<sub>2</sub> in the culture medium: Withdraw 0.1 mL from the culture and place in a small test tube. Add one drop of Nitrate Reagent A followed by one drop of Nitrate Reagent B, mixing after each addition. Pink color development occurs within



10 minutes and is directly related to amount of  $NO_2$  present. Initial tests will probably be dark magenta reflecting the 3 mM  $NO_2$  concentration of medium #480; but as culture grows and  $NO_2$  isoxidized, color will become magenta, pink and finally clear. This process can take up to 3 weeks depending on the viability of the stock.

- 4. When all  $NO_2$  has been oxidized, inoculate a 250 mL flask containing 100 mL of medium #480 with the 5.0 mL tube culture. Wrap the flask in foil to protect from light and incubate at 26°C with gentle shaking. Monitor  $NO_2$  concentration daily.
- 5. When  $NO_2$  is again depleted in the culture medium, either transfer the culture to fresh media or feed the culture by adding additional  $NO_2$ . A 10% inoculum is recommended for transfers. To feed cultures, aseptically add sterile  $NaNO_2$  solution, returning  $NO_2$  concentrations to approximately 10 mM (3.5 mL of 2%  $NaNO_2$  per 100 mL culture). Cells should be harvested or subcultured to fresh media when cultures can utilize 10 mM  $NO_2$  in less than 24 hours and cell numbers are greater than 10 cells per field at 1000X magnification.
- 6. Cells can be harvested by centrifugation. For longterm storage, 1.0 mL aliquots of concentrated cells can be frozen with glycerol as the cryoprotectant at a final concentration of 10%. Frozen vial should be stored at 70°C or below for best preservation

#### Notes

This item will not produce turbidity in broth. Growth can be detected by testing for the depletion of nitrite from the broth.

Reagent A:

Sulfanilic acid, 8 g

5N Acetic acid, 1.0 L

NO<sub>2</sub> Reagent B:

N,N-Dimethyl-1-naphthylamine, 6.0 ml

5N Acetic acid, 1.0 L

51918

5N acetic acid consists of 1 part glacial acetic acid to 2.5 parts distilled water. NO<sub>2</sub> Reagents A and B may also be purchased from Remel. Additional information on this culture is available on the ATCC® web site at www.atcc.org.

#### **Material Citation**

If use of this material results in a scientific publication, please cite the material in the following manner: *Nitrobacter* sp. (ATCC 51918)

#### References

References and other information relating to this material are available at www.atcc.org.

## Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.



#### **Disclaimers**

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

## Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.



51918

### Revision

This information on this document was last updated on 2024-10-24

### **Contact Information**

ATCC

10801 University Boulevard Manassas, VA 20110-2209

**USA** 

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor

