



Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc- Halotag

ACS-5007™

Description

Organism: *Homo sapiens*, human

Cell Type: neural progenitor cell

Tissue: Umbilical cord; CD34+ cord blood

Age: neonate

Gender: Male

Morphology: short spindle shape

Growth properties: Adherent

Disease: Normal

Cells per vial: $\geq 1.0 \times 10^6$

Volume: 1.0 mL

Storage Conditions

Product format: Frozen

Storage conditions: Vapor phase of liquid nitrogen

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

Astrocyte, oligodendrocyte, and neuron differentiation; drug screening; Quantitative measurement of early neuron differentiation.

Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag

ACS-5007

BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

Cells contain Sendai virus (SeV) DNA sequences

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Seeding density:

Post-thaw: 8.0×10^4 viable cells/cm² on Cell Basement Membrane Gel-coated dishes/

Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag

ACS-5007

Product Sheet

Subculture: 4.0×10^4 viable cells/cm² on Cell Basement Membrane Gel-coated dishes/plates

Handling Procedures

Unpacking and storage instructions:

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

Complete medium: Complete growth media for Neural Progenitor Cells (NPCs) includes DMEM: F12 (ATCC® [30-2006](#)) supplemented with the Growth Kit for Neural Progenitor Cell Expansion (ATCC® [ACS-3003](#)). To make complete NPC medium add the following components of the kit to 464 mL DMEM: F12:

- 5 mL L-Alanyl-L-Glutamine
- 5 mL Non-Essential Amino Acids
- 10 mL NPC Growth Kit Component A
- 5 mL NPC Growth Kit Component B
- 1 mL NPC Growth Kit Component C
- 10 mL NPC Growth Kit Component D

Handling Procedure:

Coat plates with Cell Basement Membrane Gel (ATCC® [ACS-3035](#)) and culture the NPCs with NPC Growth Medium (ATCC® [ACS-3003](#)).

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If, upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -80°C.

Storage at -80°C will result in loss of viability. Preparation of Cell Basement Membrane Gel (ATCC® ACS-3035™) coated plates:

1. Thaw Cell Basement Membrane Gel on ice or at 4°C
2. Prepare a 150 µg/ mL working concentration of Cell Basement Membrane Gel in cold DMEM: F-12 medium
3. Add enough Cell Basement Membrane Gel solution to cover the surface of the plate (e.g. 1 mL diluted Cell Basement Membrane Gel/well of a 12-well plate)

Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag

ACS-5007

4. Incubate for 1 hour at 37°C prior to use

Initiation of Cultures

1. Prepare complete NPC growth medium (ATCC® ACS-3003™) following the instructions in the package and pre-warm that medium as well as DMEM:F12 in a 37°C water bath for 15-30 min. If using a small volume of medium (50 mL or less), warm only the volume needed in a sterile conical tube. Avoid warming complete medium multiple times.
2. Obtain a 12-well plate with Cell Basement Membrane Gel. Aspirate the Cell Basement Membrane Gel medium and directly add 1.5 mL of the complete NPC Growth Medium per well. Place the plate in the incubator for 15 minutes to allow the medium to reach its normal pH (7.0-7.6). Four to five wells of a 6-well plate may be needed for each vial of cells thawed.
3. Transfer 9 mL of pre-warmed DMEM:F12 into a 15 mL conical tube for recovery of the NPCs from the frozen stock.
4. Remove cryovial of frozen cells from liquid nitrogen storage.
5. Thaw the cells by gently swirling in a 37°C water bath. To reduce the possibility of contamination, keep the cap out of the water. Thawing should be rapid (approximately 1 to 2 minutes). Remove the cryovial from water bath when only a few ice crystals are remaining.
6. Sterilize the cryovial with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
7. Remove cells from the vial using a P1000 micropipette and transfer cells drop-wise into the 15 mL conical tube containing 9 mL DMEM:F12.
8. Centrifuge cells at 270 x g for 5 minutes at room temperature.
9. Aspirate the supernatant and discard. Gently tap the bottom of the tube to loosen the cell pellet.
10. Add 4 mL of the complete NPC Growth Medium to the tube. Gently resuspend the pellet by pipetting up and down 3 or 4 times to make a single-cell suspension.
11. Perform cell count by a Vi-Cell Analyzer or hemocytometer. Note: Don't perform cell count by a Vi-Cell Analyzer without removal of serum-free freezing medium.
12. Seed NPCs at a seeding density of 80,000 viable cells/cm² (e.g. 0.30 x10⁶/well of a 12-well plate) onto a Cell Basement Membrane Gel-coated plate described above.
13. Incubate the plate at 37°C with 5% CO₂ overnight.

Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag

ACS-5007

14. Change medium at 100% media change rate (1 mL media/well) next day and every other day thereafter.
15. Monitor cell growth and passage the cells when they reach ~95% confluency

Note: Don't passage NPCs when the cells are <85% confluency

Subculturing procedure:

Post thaw day 1, perform a 100% medium change and remove all cells that did not attach. Perform a 100% medium change every other day thereafter. Passage the cells with diluted Accutase (50% Accutase and 50% DPBS) when they reach ~95% confluence and reseed the NPCs at 40,000 viable cells/cm² on Cell Basement Membrane Gel-coated dishes/plates.

Cryopreservation:

Post thaw day 1, perform a 100% medium change and remove all cells that did not attach. Perform a 100% medium change every other day thereafter. Passage the cells with diluted Accutase (50% Accutase and 50% DPBS) when they reach ~95% confluence and reseed the NPCs at 40,000 viable cells/cm² on CellMatrix-coated dishes/plates.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag (ATCC ACS-5007)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag

ACS-5007

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This

Neural Progenitor Cells Derived from XCL-1 MAP2p-Nanoluc-Halotag

ACS-5007

product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-12-06

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor