

**BAA-489**<sup>TM</sup>

### Description

Strain designation: JCM 10882 [DSM 13996, RKU-10] Deposited As: Thermotoga naphthophila Takahata et al.

Type strain: Yes

### **Storage Conditions**

Product format: Freeze-dried

#### Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

#### BSL<sub>1</sub>

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of Biosafety in Microbiological and Biomedical Laboratories (BMBL), U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always

used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

### Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

#### **Growth Conditions**

**Medium:** 

ATCC Medium 2332: Thermotoga petrophila medium

Temperature: 70°C
Atmosphere: Anaerobic

## Handling Procedures

- 1. Sterilize the top of the Balch tube by spraying it with 70% ethanol and then flaming the top.
- 2. If needed, exchange the gas in the test tube for 80% N<sub>2</sub> -20% CO<sub>2</sub> or 100% N<sub>2</sub>.
- 3. If the medium is pink (see discussion about resazurin) add 1 drop of reducing agent (1.5% sodium sulfide stock solution) for each one or two ml of medium. Let the medium sit at room temperature for 10 to 20 minutes, until the resazurin



becomes colorless, before inoculating.

- 4. Open the freeze-dried vial according to the enclosed instructions. Take an anaerobic (anaerobic conditions D. below) 1.0 ml syringe tipped with 22 gauge needle and withdraw 0.5 ml of medium from the Balch tube and rehydrate the freeze dried pellet. Immediately place the re-hydrated vial under a stream of sterile gas,  $80N_2$  - $20\%CO_2$  to maintain anaerobicity.
- 5. Using the same syringe, withdraw the cell suspension from the vial and transfer it to the Balch tube. Inoculate a second pre-reduced tube of medium with 0.5 ml of the rehydrated culture. Plate 0.1 ml of the inoculated culture onto a non-selective medium and incubate aerobically at  $37^{\circ}$ C. Incubate the broth tube at  $70^{\circ}$ C.
- 6. Growth should be detected in the broth within 24 to 48 hours. No growth should be detected on the aerobic plate or broth.

#### ANAEROBIC CONDITIONS: :

A. Balch tubes (available from Bellco Glass, Vineland, NJ; are specially designed for anaerobic work and use an aluminum crimp cap to hold a rubber stopper in place. Needles can easily be inserted through the stopper, and the tubes can be pressurized to 2 atm. Alternatively, serum vials may be used, or screw cap tubes with butyl rubber stoppers, in the latter case the stopper may be removed and the tube placed under a cannula system that dispenses sterile, oxygen free gas for addition of reducing agents or inoculation.

- B. Resazurin is a commonly used redox indicator that is pink when the redox potential is above 50 mv, and colorless when the redox potential is below 110 mv. i.e. highly reducing. Most strict anaerobes require this low redox potential for optimum growth.
- C. To obtain a fully reduced medium, it is necessary that the medium be anoxic and that a reducing agent be added. Common reducing agents are sodium sulfide, cysteine, dithiothreitol, and titanium citrate.
- D. Syringes can be made anaerobic by one of two methods.
  - 1. Displace the dead space in the syringe with a sterile
- 2. Displace the dead space in the syringe with a



reducing agent.

#### Notes

Once growth has been established, the culture should be transferred every 1 or 2 days. The culture will remain viable for 1 week if stored at room temperature and maintained under anaerobic conditions.

When this freeze-dried culture is incubated at elevated temperatures ( $60^{\circ}$ C or above) the cryoprotective agent precipitates. This precipitation does not interfere with growth of the organism.

Cells appear as short rod that occur singly and in pairs. When examined under phase microscopy the sheath can some times be detected.

Additional information on this culture is available on the ATCC web site at www.atcc.org.

### Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Thermotoga naphthophila* Takahata et al. (ATCC BAA-489)

#### References

References and other information relating to this material are available at www.atcc.org.

## Warranty



The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

#### **Disclaimers**

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This



product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

### Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

#### Revision

This information on this document was last updated on 2025-10-02

#### Contact Information

**ATCC** 

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor

