

Desulfovibrio bastinii Magot et al.

BAA-903[™]

Description

Strain designation: SRL4225 [DSM 16055]

Type strain: Yes

Storage Conditions

Product format: Freeze-dried

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL₁

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories* (*BMBL*), U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is

Desulfovibrio bastinii Magot et al.

BAA-903

important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Medium:

ATCC Medium 1250: Modified Barr's Medium for sulfate reducers with 2.5% NaCl

Temperature: 37°C **Atmosphere:** Anaerobic

Handling Procedures

- 1. Open the vial according to enclosed instructions.
- 2. Perform all steps under anaerobic conditions.
- 3. Aseptically transfer 0.5 ml of #1250 broth to the vial and rehydrate the pellet. Transfer the suspension back into the broth tube. Inoculate a plate of a non-selective medium such as Tryptic Soy, Nutrient, or blood agar with 0.1 ml of the cell suspension.



Desulfovibrio bastinii Magot et al. BAA-903

- 4. Seal the tube with a rubber stopper and incubate anaerobically at 37°C. Incubate the plate(s) aerobically as a purity check.
- 5. After one or two days, growth should be evident as indicated by turbidity through out the broth. Once growth has been established the culture should be transferred to fresh broth every 24 to 48 hours.
- 6. This culture is very sensitive to oxygen when initially rehydrated, therefore steps should be taken to avoid exposure to oxygen. When the culture exhibits good growth it will remain viable for up to 1 week if stored at 4°C under anaerobic condition.

ANAEROBIC CONDITIONS:

- · Tubes of media are placed under a gassing cannula system hooked to a source of oxygen free gas.
- \cdot All transfers are performed while the test tubes are on the cannula system with a gentle stream of oxygen free gas

flowing through the system.

- · As the test tubes are removed from the cannula system each is sealed with butyl rubber stopper thus maintaining the anaerobic headspace.
- \cdot 100% nitrogen or 80% nitrogen-20% carbon dioxide gas mixture is typically employed as the oxygen free gas source.

Notes

When examined microscopically, the cells appear as typical single (some pairs) comma-shaped rods that are motile.

Always use freshly prepared anaerobic medium. If there is any question about the anaerobic condition of the medium, the medium can be reduced with the addition of 1.5% cysteine

(2.0 ml per 100 ml of medium).

Desulfovibrio bastinii Magot et al. BAA-903

Other commonly used reducing agents are sodium sulfide, cysteine, dithiothreitol, and titanium citrate. Cysteine is the reducing agent of choice since it does not cause the ferrous ammonium sulfate to precipitate.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Desulfovibrio bastinii* Magot et al. (ATCC BAA-903)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Desulfovibrio bastinii Magot et al. BAA-903

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.



Desulfovibrio bastinii Magot et al.

BAA-903

Revision

This information on this document was last updated on 2025-10-03

Contact Information

ATCC

10801 University Boulevard Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor

