



# TALL-104

CRL-11386™

## Description

TALL-104 is a T lymphoblast cell line that was established from the peripheral blood of a two-year-old male child in relapse with T-ALL. Applications include immune system disorder, immunology, and immuno-oncology research.

**Organism:** *Homo sapiens*, human

**Cell Type:** T lymphoblast

**Tissue:** Peripheral blood

**Age:** 2 years

**Gender:** Male

**Morphology:** lymphoblast

**Growth properties:** Adherent and/or suspension

**Disease:** Acute lymphoblastic leukemia ALL

**Patent depository:** This material was deposited with the ATCC Patent Depository to fulfill U.S. or international patent requirements. This material may not have been produced or characterized by ATCC. As an International Depository Authority (IDA) for patent deposits, ATCC is required to complete viability testing only at time of initial deposit of patent material. Patent deposits are made available on behalf of the Depositor when the pertinent U.S. or international patent is issued, but material may not be used to infringe the patent claims.

**Patent number:**

5,272,082

**Technical information:** ATCC Product Experience does not have technical information on patent deposits that are not produced or characterized by ATCC. Additional information can be found in the corresponding patent available from the patent holder or with the U.S. and/or international patent office.

---

## Storage Conditions

**Product format:** Frozen

**Storage conditions:** Vapor phase of liquid nitrogen

---

## Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

---

## BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

---

## Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at [www.atcc.org](http://www.atcc.org).

---

## Growth Conditions

**Temperature:** 37°C

**Atmosphere:** 95% Air, 5% CO<sub>2</sub>

## Handling Procedures

### Unpacking and storage instructions:

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

**Complete medium:** The base medium for this cell line is RPMI-1640 (ATCC 30-2001).

To make the complete growth medium, add the following components to 400 mL of the base medium:

- 100 mL Fetal Bovine Serum (ATCC 30-2020) to a final concentration of 20%.
- 1 vial IL-2 IS (10 µg), premium grade (Miltenyi cat# 130-097744\*). Contents of vial can be rehydrated with 1 mL of culture medium and then added directly to the bottle of formulated medium.

**\*Note:** The biological activity of cat# 130-097-744 is at least  $5.0 \times 10^6$  units/mg but may be as high as  $9.0 \times 10^6$  units/mg per Miltenyi. This may cause the final formulation of IL-2 to be greater than 100 units/mL

**Note:** Complete culture medium has a recommended shelf life of one month when being stored at 2-8°C. Vendor does not provide stability information, but suggested material should be good for up to a month after formulation.

**Note:** IL-2 rapidly loses its potency in medium, it is important that fresh IL-2 be used. Successful growth of this cell line is very dependent upon the quality of IL-2 used in the growth medium. ATCC recommends using the highest quality IL-2 available.

### Handling Procedure:

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the

possibility of contamination, keep the O-ring and cap out of the water.

Thawing should be rapid (approximately 2 minutes).

2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. Product Sheet TALL-104 CRL-11386 [www.atcc.org](http://www.atcc.org) Page 3 of 7 operations from this point on should be carried out under strict aseptic conditions. Transfer the vial contents to a centrifuge tube containing 9.0 mL complete culture medium and spin at approximately 400 x g for 8 minutes.
4. Resuspend cell pellet with the recommended complete medium (see the lot specific information provided on the Certificate of Analysis for the culture recommended dilution ratio) and dispense into an appropriate size culture flask (recommended on the Certificate of Analysis). The recommended seeding density for CRL-11386 is  $7.0 \times 10^5$  to  $9.0 \times 10^5$  viable cells/mL. It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the complete growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).
5. Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product.

#### Comments:

CRL-11386 has historically shown a decrease in viability the first 2-3 days post thaw, the viability will begin improving as the cells remain in culture with routine media additions and replacements as outlined in the subculture restrictions.

#### Subculturing procedure:

Cultures can be maintained by addition or replacement of fresh medium. Start cultures at least  $7 \times 10^5$  cells/mL.

**Interval:** Maintain between  $4 \times 10^5$  and  $1 \times 10^6$  cells/mL.

**Medium Renewal:** Add medium as cell density increases

**Subculturing Restrictions:** The CRL-11386 cell line is extremely sensitive to overgrowth and media exhaustion. Reseeding the cell culture must be performed every 2 to 4 days. Historically, with reseeds every 2 to 4 days, it is expected to perform a full media replacement every 14-21 days. However, full fluid changes should be performed if the cell viability drops below 85% after initially recovering

from start-up above that threshold or if the media becomes too acidic before the 14 days.

It is strongly recommended to maintain cell culture vessels in an upright position during incubation. CRL-11386 tends to exhibit increased growth abilities and higher viability when the flask(s) that maintain the cells are kept upright during incubation (i.e. on the short edge of the flask). The resulting proximity of cells seemingly triggers enhanced propagation.

**Interval:** Maintain between  $3.0 \times 10^5$  to  $1.0 \times 10^6$  cells/mL

**Reagents for cryopreservation:** RPMI-1640 + 20% Gibco Knockout SR (ThermoFisher Scientific cat# 10828010) + 10% DMSO. Freeze at a cell density of at least  $15 \times 10^6$  cells/mL. Cell culture tested DMSO is available as ATCC 4-X.

---

## Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: TALL-104 (ATCC CRL-11386)

---

## References

References and other information relating to this material are available at [www.atcc.org](http://www.atcc.org).

---

## Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a

change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

---

## Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for

damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at [www.atcc.org](http://www.atcc.org).

This material is cited in a US and/or international patent and may not be used to infringe the claims. Depending on the wishes of the Depositor, ATCC may be required to inform the Depositor of the party to which the material was furnished.

---

## Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

---

## Revision

This information on this document was last updated on 2025-12-03

---

## Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: [tech@atcc.org](mailto:tech@atcc.org) or contact your local distributor