



LUHMES

CRL-2927™

Description

LUHMES is a neuronal-like cell that was isolated from the mesencephalon of a patient. This cell line was deposited by J Lotharius in 1998 and can be used in neuroscience research.

Organism: *Homo sapiens*, human

Tissue: Brain; Mesencephalon

Age: 8 weeks gestation

Morphology: neuronal

Growth properties: Adherent

Storage Conditions

Product format: Frozen

Storage conditions: Vapor phase of liquid nitrogen

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies

and procedures as well as any other applicable regulations as enforced by your local or national agencies.

Cells contain v-myc retroviral vector

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Temperature: 37°C

Atmosphere: 95% Air, 5% CO₂

Handling Procedures

Unpacking and storage instructions:

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the

cells at a temperature below -130°C , preferably in liquid nitrogen vapor, until ready for use.

Complete medium:

Complete medium: The base medium for this cell line is ATCC-formulated DMEM:F12 Medium Catalog No. 30-2006. To make the complete growth medium, add the following components to the base medium: 1% N2 supplement (Gibco-Invitrogen Cat No 17502-048) 40 ng/ml b-FGF (basic recombinant human Fibroblast Growth Factor; Gibco-Invitrogen Cat No 13256-029) added fresh at the last moment.

Example media preparation

- 495 mL DMEM/F12 medium (ATCC 30-2006)
- 5 mL N-2 Supplement (Invitrogen catalog # 17502-048; stock concentration 100X)

Note: Complete media is stable for 2 weeks after adding N-2 when stored at 2 to 8°C . Thawed N-2 should be used immediately or stored in aliquots at -5 to -20°C . Avoid additional freeze-thaw cycles.

Note: Because of limited stability, the following component should be added to an aliquot of the above culture medium fresh prior to seeding, performing fluid additions and full fluid changes. Media supplemented with the item below expires in 10 days.

- b-FGF supplement (1000 ng/mL stock): use 40 μL per mL of culture medium

Note: b-FGF stock solution has a shelf life of 3 months after the date of preparation when stored at -20°C .

Handling Procedure:

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

Note: Culture flasks should be **pre-coated** with $50\mu\text{g/mL}$ poly-L-ornithine and then with $1\mu\text{g/mL}$ human fibronectin. (See Subculturing Procedure)

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the

- possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
 3. Transfer the vial contents to a centrifuge tube containing 9.0 mL culture medium **without bFGF** and spin at approximately 125 x g for 5 to 10 minutes.
 4. Resuspend cell pellet with the recommended complete growth medium (see the lot information on Certificate of Analysis (COA) for the culture recommended dilution ratio) and dispense into a pre-coated vented 25 cm² or 75 cm² culture flask as recommended on the COA. The recommended seeding density for CRL-1720 is 4.0 x 10⁴ to 6.0 x 10⁴ viable cells/cm². It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the complete growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).
 5. Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product.

Subculturing procedure:

These cells should be recovered from cryopreservation and subcultured only on culture flasks that are sequentially **pre-coated** with 50 µg/mL poly-L-ornithine (Sigma, Cat. No. P-3655 or equivalent) and then with 1 µg/mL Human Fibronectin (Sigma, Cat. No. F-0895 or equivalent).

Note: Use flasks with vented caps for best results.

1. Add 6-15 mL freshly diluted 50 µg/mL poly-L-ornithine to a T-75 cm² flask and allow flask to incubate while rocking at room temperature at slow speed for 12 to 29 hours. Note: Use non-vented caps or parafilm tapes around the caps to avoid possible evaporation of coating solution when coating.
2. Remove and discard poly-L-ornithine solution. Rinse flask 3 times with culture grade water (Hyclone cat#SH30529.01). Discard last rinse and allow flask to air-dry uncapped and standing upright in a biological cabinet. **Note:** Over drying of coated flask, may cause less attachment after subculture.
3. Add 6-15 mL of freshly diluted 1µg/mL fibronectin and incubate for 3 hours at 37°C slowly rocking. **Note:** Use non-vented caps or parafilm tapes around the

- caps to avoid possible evaporation of coating solution when coating.”
4. Remove and discard fibronectin solution. Rinse flask 3 times with culture grade water (Hyclone cat#SH30529.01). Discard last rinse and allow flask to air-dry uncapped and standing upright in a biological cabinet. **Note:** Over drying of coated flask, may cause less attachment after subculture.
 5. Flask is ready for use when dry.

Volumes used in this protocol are for 75 cm² flasks; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes.

Notes: Subculture before cells reach 80% confluency. Warm aliquots of wash medium (growth medium without bFGF) used in Step 4, freshly made complete growth medium and freshly diluted trypsin solution to 37°C prior to use. A P1000 micropipettor should be used to resuspend cells.

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 10.0 mL Ca⁺⁺/Mg⁺⁺ free Dulbecco's phosphate-buffered saline (D-PBS).
3. Add 4.0 mL of freshly diluted, pre-warmed 0.025% Trypsin-0.1 g/L EDTA solution (see formula below) to flask and incubate for 3 minutes at 37°C. Knock the flask several times against the palm of your hand to dislodge cells. Observe flask under an inverted microscope to be sure cells have come off.
4. Add 6.0 mL of pre-warmed wash medium (see Note above) and aspirate cells by gently pipetting.
5. Transfer cell suspension to a centrifuge tube and spin at approximately 190 x g for 7 minutes. Discard supernatant and knock the tube against the palm of your hand to loosen the cell pellet. Using a 1 mL pipet, add 1.0 mL complete growth medium and pipet the pellet up and down to resuspend the cells (avoid creating foam or bubbles).
6. Add an additional 2.0 mL of complete growth medium and dissociate cells further by pipetting up and down.
7. Adjust cell concentration by adding the necessary volume of complete growth medium needed to seed new flasks. Pipet up and down to evenly resuspend cells. Cell cultures can be established between 5.0 x 10⁴ to 8.0 x 10⁴ viable cells/cm². Seed cells at subculture at a concentration between 2.0 x 10⁴ to 6.0 x 10⁴ viable cells/cm².
8. Add appropriate aliquots of the cell suspension to new pre-coated vented culture flasks.
9. Incubate cultures in 5% CO₂/95% air at 37°C.

Subcultivation ratio/Subculture seeding density: A subcultivation ratio of 1:3 to 1:10 is recommended or subculture at a seeding density of 4.0×10^4 to 8.0×10^4 viable cells/cm² is recommended. Subculture approximately every 2 to 4 days.

Medium renewal: Every 2 to 3 days

2X Trypsin-EDTA Solution: 2X Trypsin-EDTA solution is 0.05% Trypsin-0.2g/L EDTA. Before use this must be diluted 1:1 in Ca⁺⁺/Mg⁺⁺ free Dulbecco's phosphate-buffered saline (D-PBS) to 0.025% Trypsin-0.1g/l EDTA

To make 1 liter of 2X Trypsin-EDTA solution:

1. Add the following to 500 ml ddH₂O:
 - a. 8 g NaCl
 - b. 0.4 g KCL
 - c. 0.58 g NaHCO₃
 - d. 1 g Dextrose
 - e. 0.2 g EDTA
 - f. 0.5 g Trypsin (Sigma, Cat. No. T7409)
2. Bring volume up to 1 liter with ddH₂O, and pH to 7.4 with HCL.
3. Incubate at 37°C for at least 1 hour to activate the trypsin.
4. Sterile filter (0.2 µm) and make aliquots.
5. Refrigerate at 4°C overnight and then store at -20°C.
6. Before use, dilute trypsin 1:1 with Ca⁺⁺/Mg⁺⁺ free Dulbecco's phosphate-buffered saline (D-PBS) and warm to 37°C.

Reagents for cryopreservation: Complete growth medium supplemented with 20% heat inactivated fetal bovine serum and 10% (v/v) DMSO (ATCC 4-X)

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: LUHMES (ATCC CRL-2927)

References

References and other information relating to this material are available at

www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and

assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-12-18

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

LUHMES

CRL-2927

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
