



MPC 862L

CRL-3540™

Product Sheet

Description

MPC 862L is a cell line isolated from a mouse pheochromocytoma.

Organism: *Mus musculus*, mouse

Tissue: Adrenal gland

Morphology: Epithelial like and/or rounded

Disease: Pheochromocytoma

Cells per vial: 6.0×10^6

Storage Conditions

Product format: Frozen

Storage conditions: Vapor phase of liquid nitrogen

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Temperature: 37°C (35-37°C)

Atmosphere: 95% Air, 5% CO₂

Handling Procedures

Unpacking and storage instructions:

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

Complete medium:

The base medium for this cell line is ATCC-formulated RPMI-1640 Medium, Catalog No. 30-2001. To make the complete growth medium, add the following components to the base medium:

- heat-inactivated horse serum to a final concentration of 10%
- fetal bovine serum to a final concentration of 5%

Subculturing procedure: Before initiating subculture flasks must be coated with 50 µg/ mL collagen I from rat tail (ThermoFisher Scientific catalog #: A1048301 or equivalent). It is recommended to follow the vendor instructions for thin coating procedure.

It is recommended to subculture at or before 80% confluence. Cells may not reach 100% confluence and grow in clumps which will become necrotic if not passaged. Before beginning subculture procedure prepare a solution of 0.25% Gibco™ Trypsin (ThermoFisher Scientific catalog # 15050- 065) + 80 µg/mL Deoxyribonuclease I. Deoxyribonuclease I (DNase I) should be added to an aliquot of 0.25% Gibco™ Trypsin fresh prior to use.

To prepare 10 mg/mL DNase I stock solution, aseptically combine: o 100 mg Roche™ DNase I (Sigma catalog # 11284932001) o 10 mL Molecular Grade Water (ATCC catalog # 60-2450) Add 8 µL stock DNase I per 1 mL 0.25% Trypsin immediately prior to use.

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes.

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 mL of 0.25% Gibco™ Trypsin (ThermoFisher Scientific catalog # 15050-065) + 80 µg/mL Deoxyribonuclease I) to flask and observe cells under an inverted microscope until the cell layer is dispersed (usually within 5 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting.
5. Centrifuge the cell suspension at approximately 150 to 400 x *g* for 8 to 12 minutes to remove dissociation agent.
6. Resuspend the cell pellet in an appropriate amount of complete culture medium
7. If cell clumps are present then clumps should be disrupted using a 5 mL serological pipette followed by a P1000.
8. Add appropriate aliquots of the cell suspension to new culture vessels.
Cultures can be established between 1.0×10^{54} and $2.06.0 \times 10^5$ viable cells/cm².
9. Incubate cultures at 37°C.

Interval: Maintain cultures at a cell concentration between 1.0×10^{54} and 2.0×10^5

cell/cm².

Subcultivation Ratio: A subcultivation ratio of 1:2 to 1:3 is recommended

Medium Renewal: Every 2 to 3 days

Reagents for cryopreservation:

Gibco™ Recovery™ Cell Culture Freezing Media (ThermoFisher Scientific catalog # 12648-010).

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: MPC 862L (ATCC CRL-3540)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP

standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a **license from ATCC**.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-10-14

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
