



Ker-CT

CRL-4048™

Product Sheet

Description

Ker-CT is an hTERT-immortalized keratinocyte cell that was isolated from the foreskin of a male patient. This cell line was deposited by J Shay and can be used in toxicology research.

Organism: *Homo sapiens*, human

Cell Type: keratinocyte

Tissue: Skin; Foreskin

Age: neonate

Gender: Male

Morphology: epithelial

Growth properties: Adherent

Disease: Normal

Storage Conditions

Product format: Frozen

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to

Ker-CT

CRL-4048

understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

Cells contain SV40 sequences

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Temperature: 37°C

Atmosphere: 95% Air, 5% CO₂

Handling Procedures

Unpacking and storage instructions:

1. Check all containers for leakage or breakage.

Ker-CT

CRL-4048

2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

Complete medium: KGM-Gold™ BulletKit™ (Lonza 00192060). Note: Discard the GA-1000

Handling Procedure: To ensure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt.

If storage of the frozen culture is necessary upon arrival, store the vial in liquid nitrogen vapor phase and NOT at -70°C. Storage at -70°C will result in loss of viability.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All operations from this point on should be carried out under strict aseptic conditions.
3. Resuspend the cells in 5mL of complete growth medium. Count the cells and seed at recommended seeding densities. DO NOT centrifuge after thawing to remove DMSO.
4. Prepare a 25 cm² or a 75 cm² culture flask containing the recommended complete culture medium. Refer to Certificate of Analysis for recommended flask size and seeding density. Prior to the addition of vial contents, the vessel containing the growth medium should be placed in the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6) and to avoid excessive alkalinity of the medium during recovery of the cells.
5. Incubate the culture at 37°C in a suitable incubator.
6. Change to fresh medium after the cells attached, usually 6-12 hours later, to remove DMSO and FBS.

Subculturing procedure: Volumes used in this protocol are for 75 cm² flasks; proportionally reduce or increase amount of dissociation solutions for culture vessels of other sizes.

1. Remove and discard spent medium.
2. Briefly rinse with HBSS (ATCC 30-2213), 1 mL/25 cm² and discard rinse solution.
3. Add trypsin for primary cells (ATCC PCS-999-003), 1mL / 25cm². Place at 37°C for 4-6 minutes, until 90% of the cells have detached.
4. Rap flask gently to ensure cells are detached. Add 2% FBS in D-PBS, 1 mL/25cm² to neutralize trypsin.

Ker-CT

CRL-4048

5. Centrifuge cells at 250 x g for 5 min at room temperature.
6. Remove supernatant. Resuspend pellet in 6.0 to 8.0 mL Complete Growth Medium.
7. Count cells, and seed 5.0 x 10³ to 8.0 x 10³ viable cells/cm² to new culture vessels.

Medium Renewal: Every 2-3 days.

As the cells become more confluent, increase the volume of media as follows: under 25% confluence feed cells 5 mL per 25 cm², 25-45% confluence then feed cells 7.5 mL per 25 cm², over 45% confluence then feed cells 10 mL per 25 cm².

Reagents for cryopreservation: Fetal bovine serum supplemented with 10% (v/v) DMSO (ATCC 4-X)

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: Ker-CT (ATCC CRL-4048)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a

Ker-CT

CRL-4048

change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for



ATCC® Credible leads to Incredible®

www.atcc.org

damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-12-20

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
