



hTERT Neonatal Dermal Melanocytes

CRL-4064™

Description

hTERT-immortalized Neonatal Melanocytes (ATCC® CRL-4064™) were created by selecting a single cell clone from primary melanocytes that had been retrovirally transduced with the gene encoding the human telomerase catalytic subunit (hTERT). Essential melanocyte character post-transduction was confirmed by immunofluorescence staining for the melanocyte marker TRP1. The CRL-4064 cell line also retains typical melanocyte morphology as observed through phase contrast microscopy.

For-profit customers intending to use this product for non-commercial screening must include the one-time "ATCC Screening Fee" (ATCC® ACS-2103F) with their first purchase of this product.

Organism: *Homo sapiens*, human

Tissue: Skin

Age: neonate

Gender: Male

Morphology: Fibroblast-like

Growth properties: Adherent

Cells per vial: Approximately 1.5×10^6

Volume: 1.0 mL

Storage Conditions

Product format: Frozen

Storage conditions: Vapor phase of liquid nitrogen

Intended Use

hTERT Neonatal Dermal Melanocytes

CRL-4064

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submerged in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submerged in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

hTERT Neonatal Dermal Melanocytes

CRL-4064

Temperature: 37°C

Atmosphere: 95% Air, 5% CO₂

Handling Procedures

Complete medium:

The base medium for these cells is Dermal Cell Basal Medium (ATCC PCS-200-030). To make the complete medium, add the components of Adult Melanocyte Growth Kit (ATCC PCS-200-042) according to manufacturer instructions.

It's important to note that CRL-4064 requires two distinct media formulations:

- **Recovery/Thawing medium:** Dermal Cell Basal Medium (PCS-200-030) + Adult Melanocyte Growth Kit (PCS-200-042) + 30% FBS

Prepare **Recovery/Thawing** medium by adding 12 mL FBS (ATCC 30-2020) to 28 mL of the Complete medium

Note: The above media is stable for 4 weeks at 4 to 8°C.

FBS should not be heat inactivated as this can reduce or destroy serum growth factors, attachment factors and hormones.

- **Growth medium:** Dermal Cell Basal Medium (PCS-200-030) + Adult Melanocyte Growth Kit (PCS-200-042) + 0.5 µg/mL puromycin

Prepare **Growth medium** by adding 24 µL Gibco Puromycin (10 mg/mL; ThermoFisher Scientific catalog # A11138-03 or equivalent) per 480.3 mL Complete medium.

Handling Procedure:

To ensure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

1. Use 8–10 mL of prepared **Recovery/Thawing** medium and place it in a T25 flask. Equilibrate the medium in a 37°C, 5% CO₂ incubator for 30 minutes before

hTERT Neonatal Dermal Melanocytes

CRL-4064

- cell thawing.
2. Thaw the cryovial rapidly in a 37°C water bath, keeping the O-ring and cap dry. Thawing should take no more than 2 minutes.
 3. Before the last ice crystal has melted, promptly remove the vial from the water bath. This helps prevent osmotic shock and temperature stress that may compromise cell viability.
 4. Immediately disinfect the outside of the vial with 70% ethanol. From this point forward, proceed under strict aseptic conditions
 5. Transfer the vial contents into a centrifuge tube containing 1.0 mL of equilibrated thawing medium. **Do NOT pipette up and down to mix.** Rinse the vial with an additional 1.0 mL of thawing medium and add to the same tube. **Do NOT pipette up and down to mix.**
 6. Centrifuge at approximately 300 x g for 8 minutes, then remove the cryoprotectant. Resuspend cell pellet into Recovery/Thawing medium. Incubate the culture at 37°C and 5% CO₂
 7. After 24 hours, replace medium with **Growth medium** (complete medium supplemented with 0.5ug/ml puromycin) and incubate the culture at 37°C and 5% CO₂

Subculturing procedure:

Volumes used in this protocol are for 75 cm² flask; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes. Corning[®] T-75 flasks (catalog #430641) are recommended for subculturing this product.

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with PBS (ATCC 30-2200) to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 mL of Trypsin-EDTA for Primary Cells Solution to flask, ensure complete coverage of cells and remove trypsin, then incubate and observe cells until cells have detached (usually within 2 to 3 minutes).
4. Once detached, add 2.0 to 3.0 mL of a 0.05% soybean trypsin inhibitor and aspirate cells. Transfer cell suspension into a 15ml conical tube.
5. Add 4.0 to 7.0 mL of complete growth medium to flask to wash and recover residual cells, aspirate cells by gently pipetting. Transfer suspension into previous 15ml conical tube.
6. Collect cells by centrifugation at 150xg for 5min.
7. Resuspend cell pellet in 1.5 to 3.0 mL of complete medium. Add appropriate aliquots of the cell suspension to new culture vessels.

Cultures should be established from cryopreservation between 2.0 x 10⁴ and

hTERT Neonatal Dermal Melanocytes

CRL-4064

3.0 x 10⁴ viable cells/cm².

8. Incubate cultures at 37°C.

Interval: Maintain cultures at a cell concentration between 5.0 X 10³ and 5.0 X 10⁴ cell/cm².

Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:10 is recommended

Medium Renewal: 3 times per week

Reagents for cryopreservation: 100% Stem Cell Freeze Medium (ATCC ACS-3020)

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: hTERT Neonatal Dermal Melanocytes (ATCC CRL-4064)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP

standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

hTERT Neonatal Dermal Melanocytes

CRL-4064

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2026-03-28

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
