



Entamoeba histolytica Schaudinn

PRA-352™

Description

Strain designation: SAW 408 RR clone A

Deposited As: *Entamoeba histolytica* Schaudinn

Type strain: No

Storage Conditions

Product format: Frozen

Storage conditions: -80°C or colder for 1 week, vapor phase of liquid nitrogen for long-term storage

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Medium:

ATCC Medium 1171: TYGM-9 medium

Instructions for complete medium: ATCC Medium 1171 grown with mixed bacteria (Quality controlled lots of this medium are commercially available from ATCC as cat. no. PRA-1171)

Temperature: 35°C

Atmosphere: Microaerophilic

Culture system: Xenic

Handling Procedures

Storage and Culture Initiation

Frozen ampules packed in dry ice should either be thawed immediately or stored in

liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

1. One day before thawing a frozen ampule inoculate two tubes of ATCC medium 1171 with the bacterial flora only. Incubate the tubes on a 15° horizontal slant at 35°C. If the specific bacterial flora associated with this culture are not available, skip this step and proceed to step 2.
2. On the following day combine 4.1 mL ATCC medium 1171 with 0.9 mL of HIBS (heat-inactivated bovine serum) to produce 5 mL of medium enriched with 20% serum. Invert gently several times to mix. **Note:** If bacterized medium 1171 was prepared in step 1, use the bacterized medium in place of fresh medium for faster recovery/growth.
3. Remove the frozen ampule from liquid nitrogen and flame gently at the base of the cap. Remove the cap and aseptically add 0.5 mL of the serum-enriched medium prepared in step 2. Place in a 35°C water bath until thawed (2-3 min). **Note:** Manipulations of the ampule before placing in the water bath should be done as quickly as possible to avoid warming of the contents at a suboptimal rate.
4. Transfer contents of the thawed ampule to a one-dram screw-capped vial (vial holds approximately 4.0 mL).
5. Add 2.5 mL of serum-enriched medium prepared in step 2 to the vial in dropwise fashion. Tighten the cap and incubate on a 15° horizontal slant at 35°C for 2-3 hours.
6. Ice the vial for 10 minutes, then invert gently 10 times. Centrifuge the vial at 100-200 x g for 5 min.
7. Aspirate the supernatant leaving approximately 0.5 mL. **Note:** Do not aspirate the pelleted material.
8. Replace the supernatant with 3.0 mL of the bacterized medium 1171 prepared in step 1 if available; otherwise use fresh medium 1171 (i.e., not serum-enriched).
9. Incubate the vial on a 15° horizontal slant at 35°C with the cap screwed on tightly. Observe the culture daily and transfer when many trophozoites are observed (i.e., early stationary phase).

Culture maintenance:

1. Ice a test tube culture at or near peak density for 10 minutes, invert 20 times

and aseptically transfer a 0.1 and 0.3 mL aliquot to a fresh tube of ATCC medium 1171.

2. Screw cap on tightly and incubate on a 15° horizontal slant at 35°C. Transfer when many trophozoites are observed (i.e., early stationary phase).

Notes: In general, better growth can often be achieved by using media pre-inoculated with bacteria. Addition of penicillin G at 75 U/mL and streptomycin at 75 µg/mL to ATCC Medium 1171 may be necessary if the bacterial density in the culture is too high. Inoculate fresh tubes of media with bacteria at least one day prior to subcultivation and prior to addition of antibiotics, if used. Also, nutrients in ATCC medium 1171 may be diluted through use of an osmotically-balanced saline buffer such as Dulbecco's PBS (ATCC 30-2200) with the medium. Mixtures in the range of 1:1 to 3:1 (medium to buffer) usually work well to control or limit bacterial growth in cultures where other methods are not sufficient.

Reagents for cryopreservation: CPMB-5 Cryoprotective Solution

DMSO, 1.0 mL

2.5 M Sucrose, 0.8 mL

L-Cysteine/Ascorbic Acid Solution, 0.2 mL

CPMB-2 Basal Solution, 6.0 mL

HIBS, 2.0 mL

CPMB-2 Basal Solution

Yeast Extract, 60.0 g

K₂HPO₄, 1.0 g

KH₂PO₄, 0.6 g

NaCl, 2.0 g

Distilled water, 1.0 L

Autoclave for 15 minutes.

L-Cysteine/Ascorbic Acid Solution

L-Cysteine-HCL, 1.0 g

Ascorbic Acid, 0.1 g

Distilled water, 10.0 mL

Add 9.0 mL of distilled water to a 20 mL beaker and dissolve the first two components. While stirring, adjust the pH to 7.2 with 10N NaOH (approximately 0.7 mL). Adjust final volume to 10 mL with distilled water and filter sterilize. Solution

should be used soon after preparation. Discard any unused solution.

Cryopreservation:

1. Harvest cells from several cultures which are in the late logarithmic to early stationary phase of growth. Place culture vessels on ice for 10 min.
2. Invert tubes 20 times and centrifuge at 200-300 x g for 5 min.
3. While cells are centrifuging, prepare the cryoprotective solution.
 - a. Place 1.0 mL of DMSO in a 16 x 125 mm screw-capped tube and ice until solidified.
 - b. Add 0.8 mL of the 2.5 M Sucrose solution, remove from ice and invert until the DMSO is liquefied. Return to ice bath.
 - c. Add 0.2 mL of the L-Cysteine/Ascorbic Acid solution to the DMSO solution and mix.
 - d. Add 6.0 mL of the CPMB-2 Basal Solution and mix.
 - e. Add 2.0 mL HIBS (heat-inactivated bovine serum) and mix.
4. Resuspend the cell pellets and pool to a final volume of approximately 10 mL with the supernatant. Make a determination of the cell density and adjust the concentration of the cells between $5 \times 10^5/\text{mL}$ - $1 \times 10^6/\text{mL}$ using fresh medium. If the cell concentration is below $5 \times 10^5/\text{mL}$, centrifuge the cell suspension and resuspend the pellet in a volume that will yield the desired concentration.
5. After the cell concentration is adjusted, centrifuge as in step 2.
6. Remove as much supernatant as possible and determine the volume removed.
7. Resuspend the cell pellet with a volume of the cryoprotective solution equal to the volume of the supernatant removed. Invert the tube several times to obtain a uniform cell density.
8. Dispense 0.5 mL aliquots into 1.0 - 2.0 mL plastic sterile cryules (special plastic vials for cryopreservation).
9. Place vials in a controlled rate freezing unit. From room temperature cool at $-1^\circ\text{C}/\text{min}$ to -40°C . If freezing unit can compensate for the heat of fusion, maintain rate at $-1^\circ\text{C}/\text{min}$ through heat of fusion. At -40°C plunge ampules into liquid nitrogen.
10. Store ampules in a liquid nitrogen refrigerator until needed.
11. One day before thawing a frozen ampule inoculate two tubes of ATCC medium 1171 with the bacterial flora only. Incubate the tube on a 15° horizontal slant at 35°C . If the specific bacterial flora associated with this culture are not available, skip this step and proceed to step 12.
12. On the following day combine 4.1 mL of the bacterized medium 1171 prepared

in step 11 with 0.9 mL of HIBS (heat-inactivated bovine serum) to produce 5 mL of medium enriched with 20% serum. Invert gently several times to mix.

13. Remove the frozen ampule from liquid nitrogen and flame gently at the base of the cap. Remove the cap and aseptically add 0.5 mL of the serum-enriched medium prepared in step 12. Place in a 35°C water bath until thawed (2-3 min).

Note: Manipulations of the ampule before placing in the water bath should be done as quickly as possible to avoid warming of the contents at a suboptimal rate.

14. Transfer contents of the thawed ampule to a one-dram screw-capped vial (vial holds approximately 4.0 mL).
15. Add 2.5 mL of serum-enriched medium prepared in step 12 to the vial in dropwise fashion. Tighten the cap and incubate on a 15° horizontal slant at 35°C for 2-3 hours.
16. Ice the vial for 10 minutes, then invert gently 10 times. Centrifuge the vial at 100-200 x g for 5 min.
17. Aspirate the supernatant leaving approximately 0.5 mL. **Note:** Do not aspirate the pelleted material.
18. Replace the supernatant with 3.0 mL of the bacterized medium 1171 prepared in step 11.
19. Incubate the vial on a 15° horizontal slant at 35°C with the cap screwed on tightly. Observe the culture daily and transfer when many trophozoites are observed (i.e., early stationary phase).

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Entamoeba histolytica* Schaudinn (ATCC PRA-352)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This

product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2024-10-26

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor