Equid herpesvirus 1

VR-3396™

Description

Equid herpesvirus 1 strain Kentucky D is propagated in Vero cells (ATCC CCL-81). This strain was derived from ATCC VR-700 by passaging the virus two times in CRFK cells ATCC CCL-94 and three times in Vero ATCC CCL-81 cells.

Strain designation: Kentucky D
Common name: Equine abortion virus
Deposited As: Equine herpesvirus type 1

Storage Conditions

Product format: Frozen
Storage conditions: -70°C or colder

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 2

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Host: Vero (ATCC CCL-81)
Effects: cell rounding; cell sloughing; cell detachment; cell degeneration
Complete medium: EMEM (ATCC 30-2003) + 2% FBS (ATCC 30-2020)
Temperature: 37°C
Recommendations for infection: Plate cells 24-48 hours prior to infection and infect when cultures are 80-90% confluent. Remove medium and inoculate with a small volume of virus (e.g. 1 mL per 25 cm^2) diluted to provide an optimal MOI (e.g. 0.01). Adsorb 1-2 hours at 37°C in a humidified 5% CO_2 atmosphere. End adsorption by adding virus growth medium.
Incubation: 3-5 days at 37°C in a humidified 5% CO_2 atmosphere, until CPE is progressed through 90-99% of the monolayer.

Equid herpesvirus 1

VR-3396

Notes

ATCC VR-3396 was derived from ATCC VR-700 by passaging the virus two times in CRFK cells ATCC CCL-94 and three times in Vero ATCC CCL-81 cells.

Key Abbreviations: °C, Degrees Celsius; CO_2, Carbon dioxide; EMEM, Eagle's Minimum Essential Medium; FBS, Fetal bovine serum; MOI, Multiplicity of infection

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: Equid herpesvirus 1 (ATCC VR-3396)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

Equid herpesvirus 1

VR-3396

Revision

This information on this document was last updated on 2025-03-10

Contact Information

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor