

Chlamydia pecorum Fukushi and Hirai

VR-628[™]

Description

Strain designation: Sporadic Bovine Encephalomyelitis [strain E58 (McNutt)]

Deposited As: Chlamydia psittaci (Lillie) Page

Type strain: Yes

Storage Conditions

Product format: Frozen

Storage conditions: -70°C or colder

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL₂

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of Biosafety in Microbiological and Biomedical Laboratories (BMBL), U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.



Chlamydia pecorum Fukushi and Hirai

VR-628

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Host: McCoy [McCoy B] (ATCC CRL-1696)

Effects: CPE; cytoplasmic inclusions; death of host animal; hemorrhage; hyperemia of

yolk sac

Complete medium:

DMEM (ATCC $^{\circ}$ 30-2002 $^{\circ}$) + 10% prescreened FBS + 10 mM HEPES + 2 μ g/mL

Cycloheximide (Sigma C-4859 Ready-Made)

Temperature: 37°C

Atmosphere: 95% Air, 5% CO₂

Recommendations for infection: For best results cells should be 24 to 48 hours old

and 90-100% confluent as the cylcoheximide will prevent further growth.

Incubation: 48-72 hours

Handling Procedures



Chlamydia pecorum Fukushi and Hirai VR-628

Mycoplasma contamination: Not detected

Notes

FBS used to culture *Chlamydia* must be prescreened to verify that the serum does not contain antibodies to *Chlamydia* or other factors that would interfere with growth.

Highly lethal and infectious for guinea pigs as well as chicken embryos. Non-infectious for mice, pigeons, sparrows, parakeets, turkeys and lambs. Growth rate markedly increases with temperature, optimal temperature for growth in chicken embryos is 39°C. Suggested protocol for propagation in TC: Add glass beads and vortex preparation to disrupt cells. Infect monolayer with disrupted material. Centrifuge at 3000 x rpm (750 x g) for 1 hour. Feed with fresh growth medium containing FBS prescreened for Chlamydia antibodies and 1-5mg/mL cycloheximide. Incubate at 37°C for 48 to 72 hours, a 5% $\rm CO_2$ in air atmosphere is recommended. Biosafety level 3 facilities and practices are indicated for activities with high potential for droplet or aerosol production.

Next-generation sequencing (NGS) at ATCC on the McCoy cell line (ATCC[®] CRL-1696™) used as the host has shown the presence of Mus Musculus mobilized endogenous polytropic provirus and Murine leukemia virus.

Key Abbreviations: °C, Degrees Celsius; CO₂, Carbon dioxide; DMEM, Dulbecco's Modified Eagle's Medium; FA, Fluorescent antibody assay; FBS, Fetal bovine serum; g, Acceleration of gravity; HEPES, N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid)

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Chlamydia pecorum* Fukushi and Hirai (ATCC VR-628)

References



Chlamydia pecorum Fukushi and Hirai VR-628

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

Chlamydia pecorum Fukushi and Hirai VR-628

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2025-03-24

Contact Information

ATCC 10801 University Boulevard Manassas, VA 20110-2209 USA



Chlamydia pecorum Fukushi and Hirai

VR-628

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor

