



Lysinibacillus sp. bacteriophage G

43725-B1™

Description

Strain designation: G [HER 276]

Deposited As: G

Storage Conditions

Product format: Frozen

Storage conditions: -80°C or colder

Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

ATCC highly recommends that appropriate personal protective equipment is always

used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

Growth Conditions

Host: *Lysinibacillus* sp. PGH [HER 1276] (ATCC 43725)

Medium:

ATCC Medium 1652: Beef Extract Broth

Temperature: 30°C

Atmosphere: Aerobic

Handling Procedures

1. Follow general procedures given below for phage propagation.
2. *Lysinibacillus* sp. strain PGH (ATCC[®] 43725™) is recommended as the propagation host.

GENERAL PROCEDURES FOR THE PROPAGATION OF BACTERIOPHAGE

To recover phage from freeze-dried or thawed frozen vial:

1. Prepare an actively growing broth culture of the recommended host strain 18-24 hours before opening the phage specimen. The host should be in early log phase.
2. Melt several tubes of soft agar (0.5% agar) and maintain at 45-55°C until ready to use.
3. Prewarm plates of the recommended medium in an incubator for approximately 15-30 minutes prior to the plating of the bacteriophage.
4. Add 100µl of the host culture to a pre-melted 0.5% agar tube and pipette gently to evenly mix, avoiding the creation of bubbles.
 - a. Overlay the agar plate surface with 2.5 -3 mL of melted 0.5% agar containing the host. Allow overlay to harden.
5. Perform the broth serial dilution in quadruplicate using a 96-well plate.
 - a. Dispense 90 µL of broth medium into each well.
 - b. Add 10 µL of phage lysate to each of the four 10⁻¹ wells. Mix thoroughly by pipetting up and down at least 15 times.
 - c. Transfer 10 µL from the 10⁻¹ wells to the 10⁻² wells and mix. Continue the serial dilution until at least 10⁻⁸.
 - d. Using a multichannel pipette, spot 2 µL of each dilution on the agar overlay. Allow to dry before moving the plates to the appropriate incubator.
6. Incubate plates agar side down at the appropriate temperature. Incubation times will vary depending on the phage. Most only need 12-18 hours. Incubating for too long can make the plaques very difficult to read.

To propagate phage:

1. Open the host organism according to the information on the product sheet.
2. Pick one colony from the isolation plate and homogenize in 5 mL of the appropriate broth.
 - a. Note: It may be necessary to inoculate a larger volume or more test tubes based on the volume needed for the next amplification.
3. Incubate at the appropriate temperature in a shaking incubator until it reaches OD₆₀₀ of 0.1 – 0.3.
4. Thaw or rehydrate the bacteriophage vial. Use 0.5 mL of the appropriate broth to rehydrate freeze-dried material.
 - a. Infect each 5 mL culture with 100 µL of the bacteriophage and shake at the appropriate temperature overnight. Prepare a fresh subculture of the host.

- b. Centrifuge phage culture at 4000 x g for 10 minutes.
- c. Filter the lysate with a 0.2 µm PES sterile filter and store the filtrate at 4°C.
- d. Perform a spot titer as per the instructions above.
- e. They may also be frozen with or without cryoprotectant. If available, liquid nitrogen storage is the best method for long term storage. Most phage can also be freeze-dried. We use double-strength skim milk mixed half-and-half with the filtrate.

NOTE: Broth propagation methods, such as this one, are successful with most phage.

Notes

Resistant colonies may be seen in areas of clear lysis.

In the event plaques aren't seen, 2M MgCl₂ and 2M CaCl₂ may be added to the broth used throughout the propagation procedure to help increase phage aggregation potential.

Plaques are 0.1 mm in diameter and are hazy.

We have been unsuccessful at either freezing or freeze-drying this phage. It will remain viable at 4°C for an indefinite length of time. Each vial contains 0.5 mL of crude phage lysate. The titer when last checked was >1 x 10⁸ pfu/mL.

Largest phage known, unique morphology (spring-like spiral around sheath). Contains more DNA than any other virus (500Md = 750 kb); DNA glucosylated.

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Lysinibacillus* sp. bacteriophage G (ATCC 43725-B1)

References

References and other information relating to this material are available at www.atcc.org.

Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility

of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

Copyright and Trademark Information

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

Revision

This information on this document was last updated on 2026-05-29

Contact Information

ATCC

10801 University Boulevard

***Lysinibacillus* sp. bacteriophage G**

43725-B1

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

Worldwide telephone: +1-703-365-2700

Email: tech@atcc.org or contact your local distributor
