



# *Thraustochytrium caudivorum*

PRA-210™

## Description

**Strain designation:** S4

**Type strain:** No

---

## Storage Conditions

**Product format:** Frozen

**Storage conditions:** -80°C or colder for 1 week, vapor phase of liquid nitrogen for long-term storage

---

## Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

---

## BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies and procedures as well as any other applicable regulations as enforced by your local or national agencies.

## *Thraustochytrium caudivorum*

PRA-210

ATCC highly recommends that appropriate personal protective equipment is always used when handling vials. For cultures that require storage in liquid nitrogen, it is important to note that some vials may leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vial exploding or blowing off its cap with dangerous force creating flying debris. Unless necessary, ATCC recommends that these cultures be stored in the vapor phase of liquid nitrogen rather than submersed in liquid nitrogen.

---

### Certificate of Analysis

For batch-specific test results, refer to the applicable certificate of analysis that can be found at [www.atcc.org](http://www.atcc.org).

---

### Growth Conditions

**Medium:**

ATCC Medium 2673: Thraustochytrid medium

ATCC Medium 790: By+ medium

**Instructions for complete medium:**

**Media:** ATCC Medium 2673

**Alternate Media:** ATCC medium 790

**Temperature:** 25°C

**Culture system:** Axenic

**Incubation:** Interval: monthly

---

### Handling Procedures

**Storage and Culture Initiation**

Frozen ampules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampoules may be stored at or below  $-70^{\circ}\text{C}$  for approximately one week. **Do not under any circumstance store frozen ampoules at refrigerator freezer temperatures (generally  $-20^{\circ}\text{C}$ ).** Storage of frozen material at this temperature will result in the death of the culture.

1. To thaw a frozen ampule, place it in a  $35^{\circ}\text{C}$  water bath such that the lip of the ampule remains above the water line. Thawing time is approximately 2 to 3 minutes. Do not agitate the ampule. Do not leave ampule in water bath after thawed.
2. Immediately after thawing, aseptically transfer contents to a T-25 tissue culture flask or 16 x 125 mm test tube containing 5 ml ATCC Medium 2673.
3. Screw the cap on tightly and incubate the tube or flask at  $25^{\circ}\text{C}$ .

**Culture maintenance:**

1. When the culture is at or near peak density, vigorously agitate the culture.
2. Transfer approximately 0.25 ml to a fresh tube or flask containing 5 ml of fresh ATCC medium 2673.
3. Screw the caps on tightly and incubate at  $25^{\circ}\text{C}$  (incubate test tubes at a  $15^{\circ}$  horizontal slant).
4. Thraustochytrids will eventually form an almost continuous sheet of cells on the bottom surface of the flask or test tube, with other cells floating in the liquid column and collecting at the fluid meniscus.
5. Repeat steps 1-3 at monthly intervals.

**Cryopreservation:**

1. To achieve the best results set up cultures with several different inocula (e.g. 0.25 ml, 0.5 ml, 1.0 ml). Harvest cultures and pool when the culture that received the lowest inoculum is at or near peak density.
2. If the cell concentration exceeds the required level do not centrifuge, but adjust the concentration to between  $2 \times 10^6$  and  $2 \times 10^7$  cells/ml with fresh medium. If the concentration is too low, centrifuge at  $600 \times g$  for 5 min and resuspend the pellet in the volume of fresh medium required to yield the desired concentration.
3. While cells are centrifuging prepare a 20% (v/v) solution of sterile DMSO as follows: Add the required volume of DMSO to a glass screw-capped test tube

## *Thraustochytrium caudivorum*

PRA-210

- and place it in an ice bath. Allow the DMSO to solidify. Add the required volume of refrigerated medium. Dissolve the DMSO by inverting the tube several times. \*NOTE: If the DMSO solution is not prepared on ice, an exothermic reaction will occur that may precipitate certain components of the medium.
4. Mix the cell preparation and the DMSO in equal portions. Thus, the final concentration will be between  $10^6$  and  $10^7$  cells/ml and 10% (v/v) DMSO. The time from the mixing of the cell preparation and DMSO stock solution to the start of the freezing process should be no less than 15 min and no longer than 30 min.
  5. Dispense in 0.5 ml aliquots into 1.0 - 2.0 ml sterile plastic screw-capped cryules (special plastic vials for cryopreservation).
  6. Place the vials in a controlled rate freezing unit. From room temperature cool at  $-1^{\circ}\text{C}/\text{min}$  to  $-40^{\circ}\text{C}$ . If the freezing unit can compensate for the heat of fusion, maintain rate at  $-1^{\circ}\text{C}/\text{min}$  through the heat of fusion. At  $-40^{\circ}\text{C}$  plunge into liquid nitrogen. Alternatively, place the vials in a Nalgene  $1^{\circ}\text{C}$  freezing apparatus. Place the apparatus at  $-80^{\circ}\text{C}$  for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately  $-1^{\circ}\text{C}/\text{min}$ .)
  7. The frozen preparations are stored in either the vapor or liquid phase of a nitrogen freezer.
  8. To establish a culture from the frozen state place an ampule in a water bath set at  $35^{\circ}\text{C}$  (2-3 min). Immerse the vial just sufficient to cover the frozen material. Do not agitate the vial.
  9. Immediately after thawing, aseptically remove the contents of the ampule and inoculate into 5 ml of fresh ATCC medium 2673 in a T-25 tissue culture flask or 16 x 125 mm screw-capped test tube. Incubate at  $25^{\circ}\text{C}$ .

---

### Material Citation

If use of this material results in a scientific publication, please cite the material in the following manner: *Thraustochytrium caudivorum* (ATCC PRA-210)

---

### References

References and other information relating to this material are available at

[www.atcc.org](http://www.atcc.org).

---

## Warranty

The product is provided 'AS IS' and the viability of ATCC® products is warranted for 30 days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

---

## Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a [license from ATCC](#).

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and

## ***Thraustochytrium caudivorum***

PRA-210

assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at [www.atcc.org](http://www.atcc.org).

---

### **Copyright and Trademark Information**

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

---

### **Revision**

This information on this document was last updated on 2024-10-26

---

### **Contact Information**

ATCC

10801 University Boulevard

Manassas, VA 20110-2209

USA

US telephone: 800-638-6597

***Thraustochytrium caudivorum***

PRA-210

Worldwide telephone: +1-703-365-2700

Email: [tech@atcc.org](mailto:tech@atcc.org) or contact your local distributor

---