

**ATCC medium: 1729 *Ruminococcus pasteurii* medium**

KH <sub>2</sub> PO <sub>4</sub> .....	0.2 g
NH <sub>4</sub> Cl .....	0.25 g
NaCl.....	1.0 g
MgCl <sub>2</sub> . 6H <sub>2</sub> O .....	0.4 g
KCl.....	0.5 g
CaCl <sub>2</sub> . 2H <sub>2</sub> O .....	0.15 g
Trace Elements Solution SL-7 (see below).....	1.0 ml
Resazurin.....	1.0 mg
NaHCO <sub>3</sub> .....	2.5 g
Sodium tartrate.....	2.0 g
Na <sub>2</sub> S . 9H <sub>2</sub> O .....	0.36 g
Distilled water.....	1.0 L

Adjust medium for final pH 7.2. Boil medium without sodium bicarbonate, trace elements or sodium sulfide under 80% N<sub>2</sub>, 20% CO<sub>2</sub>; dispense and tube with the same gas phase. Cool and aseptically add the filter-sterilized sodium bicarbonate and trace elements. Aseptically add the sodium sulfide which has been separately autoclaved under N<sub>2</sub>.

Trace Element Solution SL-7:

Hydrochloric acid, 25%.....	10.0 ml
FeCl <sub>2</sub> . 4H <sub>2</sub> O .....	1.5 g
CoCl <sub>2</sub> . 6H <sub>2</sub> O .....	190.0 mg
MnCl <sub>2</sub> . 4H <sub>2</sub> O .....	100.0 mg
ZnCl <sub>2</sub> .....	70.0 mg
H <sub>3</sub> BO <sub>3</sub> .....	62.0 mg
Na <sub>2</sub> MoO <sub>4</sub> . 2H <sub>2</sub> O.....	36.0 mg
NiCl <sub>2</sub> . 6H <sub>2</sub> O .....	24.0 mg
CuCl <sub>2</sub> . 2H <sub>2</sub> O .....	17.0 mg
Distilled water.....	1.0 L

Dissolve the FeCl<sub>2</sub> . 4H<sub>2</sub>O in the concentrated HCl, then dilute. Use 1.0 ml/L of medium.