



Product Sheet

Chlorella vulgaris (ATCC® 30821™)

Please read this FIRST



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Chlorella vulgaris* (ATCC® 30821™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Deposited Name: *Chlorella vulgaris* Beijerinck

Depositor: MP Starr

Isolation:

reservoir, Ukraine, USSR, 1966(?) Host for Vampirovibrio chlorellavorus ATCC 29753

Notes

Additional information on this culture is available on the ATCC web site at www.atcc.org.

While every effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

ATCC recommends that individuals contemplating commercial use of any culture first contact the originating investigator to negotiate an agreement. Third party distribution of this culture is discouraged, since this practice has resulted in the unintentional spreading of contaminated cultures.

Propagation

Growth Conditions

Temperature: 25.0°C

Duration: axenic; under 150 foot candles

Medium

ATCC® Medium 5: Sporulation agar

Instructions for Complete Medium

ATCC Medium 5

Protocols

Frozen ampules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

1. To thaw a frozen ampule, place in a 35°C water bath, until thawed (2-3 min). Immerse the ampule just sufficient to cover the frozen material. Do not agitate the ampule.
2. Immediately after thawing, aseptically transfer contents to a screw-capped borosilicate test tube containing ATCC Medium 5. Incubate the tube on a 15° horizontal slant at 50-100 μEinsteins/m²/s irradiance at 25°C with the cap loosened one half turn. Maintain under a 14/10 h light-dark photoperiod and subculture every 14-21 days.

Culture Maintenance

1. For a slant culture, transfer cells with an inoculating loop to a tube of fresh agar medium from a growing culture at or near peak density. For a broth culture, inoculate a tube of fresh broth medium with 0.1 ml from a growing culture at or near peak density.
2. Incubate at 50-100 μEinsteins/m²/s irradiance at 25° C with the cap loosened one half turn. Maintain under a 14/10 h light-dark photoperiod and subculture every 14-21 days.

Cryopreservation

1. Harvest cells from a culture which is at or near peak density by adding 3.0 ml fresh ATCC medium 5 broth to the slant and washing cells into suspension.
2. Adjust the concentration of cells to 2×10^5 - 2×10^7 / ml with fresh broth medium, then dilute to half this concentration by adding an equal amount of a 10% (v/v) sterile methanol solution in fresh ATCC medium 5 broth.
3. Dispense in 0.5 ml aliquots into 1.0 - 2.0 ml sterile plastic screw-capped cryules (special plastic vials for cryopreservation). The time from the mixing of the cell preparation and methanol stock solution to the beginning of the freezing process should be no less than 5 min and no greater than 15 min.
4. Place vials in a controlled rate freezing unit. From room temperature cool at -1°C/min to -40°C. If freezing unit can compensate for the heat of fusion, maintain rate at -1 C/min through heat of fusion. At -40°C plunge



Product Sheet

Chlorella vulgaris (ATCC® 30821™)

Please read this FIRST



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Chlorella vulgaris* (ATCC® 30821™)

- ampules into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately -1°C/min.)
- The frozen preparations should be stored in either the vapor or liquid phase of a nitrogen refrigerator. Frozen preparations stored below -130°C are stable indefinitely. Those stored at temperatures above -130°C are progressively less stable as the storage temperature is elevated. Vials can be stored between -80 and -70°C for no longer than one week.
 - To establish a culture from the frozen state place an ampule in a water bath set at 35°C. Immerse the vial to a level just above the surface of the frozen material. Do not agitate the vial.
 - Immediately after thawing, do not leave in the water bath, aseptically remove the contents of the ampule and add to a centrifuge tube containing 5 ml of ATCC medium 5 without agar. Centrifuge at 300 x g for 5 min.
 - Remove most of the supernatant (=methanol, which can inhibit growth) and then resuspend the pellet. Transfer the culture to a fresh tube of ATCC medium 5 and incubate on a 15° horizontal slant at 50-100 μ Einsteins/m²/s irradiance at 25°C with the cap loosened one half turn. Maintain under a 14/10 h light-dark photoperiod and subculture every 14-21 days.



References

References and other information relating to this product are available online at www.atcc.org.



Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/26]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor