



Product Sheet

# *Tetrahymena nipissingi* (ATCC® 30837™)

Please read this **FIRST**



## Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

## Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Tetrahymena nipissingi* (ATCC® 30837™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

## Description

**Strain Designation:** X2-AM  
**Deposited Name:** *Tetrahymena nipissingi* Nyberg  
**Depositor:** DL Nanney, EM Simon  
**Isolation:**  
derived from ATCC 27796

## Propagation

**Growth Conditions**  
**Temperature:** 25.0°C  
Duration: axenic

**Medium**  
ATCC® Medium 357: Tetrahymena medium

## Instructions for Complete Medium

ATCC Medium 357 is used for short-term cultivation. (ATCC medium 1034 can also be used for short-term cultivation and is available in a freeze-dried format from ATCC; contact sales for details). ATCC Medium 383 is used for long-term cultivation.

## Cryopreservation

### RM-9 Media for cryopreservation of *Tetrahymena*

Proteose Peptone (Difco 0120)	5.0 g
Tryptone	5.0 g
K <sub>2</sub> HPO <sub>4</sub>	0.2 g
Glucose	1.0 g
Liver extract	0.1 g
Glass distilled water	1.0 L

Dissolve components in glass distilled H<sub>2</sub>O and autoclave.

### Dryls Salt Solution

0.1 M NaH <sub>2</sub> PO <sub>4</sub> · 3H <sub>2</sub> O	10.0 ml
0.1 M Na <sub>2</sub> HPO <sub>4</sub> · 7H <sub>2</sub> O	10.0 ml
0.1 M Sodium citrate · 2H <sub>2</sub> O	15.0 ml
0.1 M CaCl <sub>2</sub> · 2H <sub>2</sub> O	15.0 ml
Distilled water	950.0 ml

Add the first 3 components to the distilled H<sub>2</sub>O and mix thoroughly.

Add the CaCl<sub>2</sub> solution and mix thoroughly.

(Adding the solutions in the order indicated will avoid the precipitation of Ca salts.)

1. Transfer *tetrahymena* from usual growth medium to RM-9 medium and allow to grow to near peak density.
2. Harvest cells from a culture by centrifugation at 300 x g for 2 min.
3. Adjust concentration of cells to 2 x 10<sup>6</sup>/ml in fresh medium.
4. While cells are centrifuging, prepare a 22% (v/v) sterile solution of sterile DMSO in fresh medium.
  - a) Add 2.2 ml of DMSO to an ice cold 20 x 150 mm screw-capped test tube;
  - b) Place the tube on ice and allow the DMSO to solidify (~5 min) and then add 7.8 ml of ice cold medium;
  - c) Invert several times to dissolve the DMSO;
  - d) Allow to warm to room temperature.
5. Add a volume of the DMSO solution equal to the cell suspension volume but add in 3 equal aliquots at 2 min intervals. Thus, the final concentration of the preparation will equal 11% (v/v) DMSO and 10<sup>6</sup> cells /ml.
6. Dispense in 0.5 ml aliquots into 1.0 - 2.0 ml sterile plastic screw-capped cryovials (special plastic vials for cryopreservation).
7. Place the ampules in a controlled rate freezing unit. The cooling cycle should be initiated no less than 15 min and no longer than 60 min after the addition of the DMSO to the cell preparation. From room temperature cool at -1°C/min to -40°C. If freezing unit can compensate for the heat of fusion, maintain rate at -1°C/min through heat of fusion. At -50°C ampules are plunged into liquid



Product Sheet

***Tetrahymena nipissingi***  
**(ATCC® 30837™)**

**Please read this FIRST**



**Intended Use**

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

**Citation of Strain**

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Tetrahymena nipissingi* (ATCC® 30837™)

nitrogen.

8. Store in the vapor or liquid phase of a nitrogen refrigerator.
9. To establish a culture from the frozen state aseptically add 0.5 ml sterile Dryl's Salt Solution to an ampule. Immediately place the ampule in a 35°C water bath, until thawed (2-3 min). Immerse the ampule just sufficient to cover the frozen material. Do not agitate the ampule.
10. Immediately after thawing, aseptically remove the contents of the ampule and inoculate into 5.0 ml of fresh medium in a 16 x 125 mm screw-capped test tube with a slightly loosened cap. Incubate at 25°C.

**CRYOPRESERVATION:**

**Alternative Thawing Procedure**

1. Aseptically add 0.5 ml of sterile modified PYNFH medium (ATCC Medium 1034) containing 8% (w/v) sucrose to the ampule. Immediately, place in a 35°C water bath, until thawed. Immerse the ampule just sufficient to cover the frozen material. Do not agitate the ampule.
2. Immediately after thawing, aseptically remove the contents of the ampule and gently add the material to the edge of a 20 x 100 mm petri plate containing ATCC Medium 919 (non-nutrient agar) and position on a 15 degree slant. The cell suspension will pool at the edge of the plate.
3. Continue to double the volume of the cell suspension at 10 minute intervals by adding ATCC medium 1034) containing 4% sucrose (w/v). When the volume reaches 16.0 ml place the plate in horizontal position and incubate at 25°C.
4. On the following day, gently remove the cell suspension for the plate and transfer to a T-25 tissue culture flask. Note the volume of the suspension and add a volume of fresh medium containing 4% sucrose equal to the volume of the cell suspension. Incubate the culture at 25°C.
5. After culture has been established subculture into fresh normal medium without sucrose.

**References**

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

**Biosafety Level: 1**

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

**ATCC Warranty**

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

**Disclaimers**

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).  
© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/06]

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor