



Product Sheet

Dunaliella bardawil (ATCC® 30861™)

Please read this FIRST



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Dunaliella bardawil* (ATCC® 30861™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Deposited Name: *Dunaliella bardawil* Amotz and Avron
Depositor: A Ben-Amotz
Isolation:
salt pond near Bardawil Lagoon, North Sinai, Israel, 1976

Propagation

Growth Conditions
Max Temperature: 25.0°C
Min Temperature: 20.0°C

Medium
ATCC® Medium 1174: DA medium

Instructions for Complete Medium
ATCC Medium 1174

Culture Maintenance

1. Inoculate a test tube containing 5 ml of fresh medium with 0.1 ml from a growing culture at or near peak density.
2. Incubate at 50-100 μ Einsteins/m²/s irradiance at 25°C. Maintain under a 14/10 h light-dark photoperiod.

Cryopreservation

1. Prepare a 5% (v/v) sterile methanol solution in fresh ATCC medium 1174 without agar.
2. Harvest cells from a culture which is at or near peak density by adding 3.0 ml methanol solution to the slant and washing cells into suspension.
3. Adjust the concentration of cells to 2 x 10⁹/ml with fresh methanol solution.
4. Dispense in 0.5 ml aliquots into 1.0 - 2.0 ml sterile plastic screw-capped cryules (special plastic vials for cryopreservation). The time from mixing of the cell preparation and the methanol solution, before the cooling cycle begins, should be no greater than 15 min.
6. Place vials in a controlled rate freezing unit. From room temperature cool at -1°C/min to -40°C. If freezing unit can compensate for the heat of fusion, maintain rate at -1 C/min through heat of fusion. At -40°C plunge ampules into liquid nitrogen. Alternatively, place the vials in a Nalgene 1°C freezing apparatus. Place the apparatus at -80°C for 1.5 to 2 hours and then plunge ampules into liquid nitrogen. (The cooling rate in this apparatus is approximately -1°C/min.)
7. Store ampules in either the vapor or liquid phase of a nitrogen refrigerator. Frozen preparations stored below
-130°C are stable indefinitely. Those stored at temperatures above -130°C are progressively less stable as the storage temperature is elevated. Shelf life must be determined empirically for storage temperatures above -130°C.
8. To establish a culture from the frozen state place an ampule in a water bath set at 35°C. Immerse the vial to a level just above the surface of the frozen material. Do not agitate the vial.
9. Immediately after thawing, do not leave in the water bath, aseptically remove the contents of the ampule and add to a centrifuge tube containing 5 ml of ATCC medium 1174 without agar. Centrifuge at 300 x g for 5 min.
10. Remove most of the supernatant and then resuspend the pellet. Place culture on ATCC medium 1174 slants and incubate on a 15° horizontal slant at 50-100 μ Einsteins/m²/s irradiance at 25°C. Maintain under a 14/10 h light-dark photoperiod.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1


Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S.



Product Sheet

Dunaliella bardawil **(ATCC® 30861™)**

Please read this **FIRST**



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Dunaliella bardawil* (ATCC® 30861™)

Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/26]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor