



Product Sheet

# *Neisseria meningitidis* (ATCC® 35561™)

Please read this **FIRST**



## Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

## Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Neisseria meningitidis* (ATCC® 35561™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

## Description

**Designation:** M-112 [BO-6]

**Deposited Name:** *Neisseria meningitidis* (Albrecht and Ghon) Murray

**Antigenic Properties:** Serogroup Y

## Propagation

### Medium

ATCC® Medium 814: GC Agar/Broth Medium

### Growth Conditions

**Temperature:** 37°C

**Atmosphere:** 5% CO<sub>2</sub>

### Propagation Procedure

1. Open vial according to enclosed instructions.
2. Using a single tube of #814 broth (5 to 6 mL), withdraw approximately 0.5 to 1.0 mL with a Pasteur or 1.0 mL pipette. Rehydrate the entire pellet.
3. Aseptically transfer this aliquot back into the broth tube. Mix well.
4. Use 0.2 mL of the suspension to inoculate a #814 slant, and/or 0.1 mL for a plate.
5. Incubate all tubes and plate at 37°C with 5% CO<sub>2</sub>. Growth should occur in 24/48 hours. Assure that screw caps are loose during incubation period.

## Notes

After 48 hours, growth is evident by entire, smooth, glistening, and low convex colonies on the agar surface. Brain Heart Infusion (BD 237500) can be used to rehydrate vial contents if #814 broth is not available, but it will not support growth as well as the GC broth. If a clear broth culture is desired, use HTM (Haemophilus Test Broth). To make HTM, a fresh hematin stock solution is prepared by dissolving 50 mg of hematin powder in 100 ml of 0.01 M NaOH with heat and stirring until the powder is dissolved thoroughly. Thirty milliliters of the hematin stock solution is added to 1.0 L of Mueller Hinton Broth with 5g of yeast extract. After autoclaving and cooling, aseptically add 3 ml of a NAD (nicotinamide adenine dinucleotide) stock solution (50 mg of NAD dissolved in 10 ml of distilled water; filter sterilized).

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).

## References

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

## Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

## ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

## Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been



Product Sheet

## ***Neisseria meningitidis*** **(ATCC® 35561™)**

confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.


Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/26]

---

### **Please read this FIRST**



---

### **Intended Use**

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

---

### **Citation of Strain**

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Neisseria meningitidis* (ATCC® 35561™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor