




Product Sheet

Ectothiorhodospira


shaposhnikovii (ATCC®)

43036™)

Please read this FIRST



Storage Temp.
Frozen: -80°C or colder
Freeze-Dried: 2°C to 8°C
Live Culture: See Propagation Section



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Ectothiorhodospira shaposhnikovii* (ATCC® 43036™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Designation: Imhoff beta 1 [BN 9512, DSM 2111]

Deposited Name: *Ectothiorhodospira vacuolata* Imhoff et al.

Propagation

Medium

ATCC® Medium 1448: Modified *Ectothiorhodospira* medium

Growth Conditions

Temperature: 26°C

Atmosphere: Anaerobic, under a tungsten lamp

Propagation Procedure

1. Open vial according to enclosed instructions. This organism can tolerate brief exposure to oxygen, so it may be opened under aerobic conditions. Aseptically transfer 0.5 mL of medium #1448 to the vial and rehydrate the pellet. Transfer this suspension back into a single, small tube (filled to capacity) of #1448 broth. Plate 0.1 mL of the culture on any non-selective media and incubate aerobically at 26°C. Seal the tube with a screw cap, and incubate at 26°C (room temperature).
2. After four to seven days, growth is evident by turbidity. When examined microscopically, the cells appear as vibroid rods, in singles and pairs, some small chains. Once growth has been detected, the culture should be transferred to fresh broth. Subsequent growth is detected within 48 to 72 hours.
3. For best results the medium should be pre-reduced, this can best be accomplished by:
 - a. Steaming the medium, then filling the test tube to capacity and sealing with a screw cap while the medium is still hot.
 - b. Adding 5 to 6 drops of cysteine (3% stock concentration) or sodium sulfide (1.5% stock concentration) to each 5 to 6 mL of medium used. After the reducing agent (cysteine or sodium sulfide) has been added fill the test tube to capacity and seal with a screw cap. Let the test tube sit at room temperature for 30 minutes before inoculating.
4. Note that once growth has been established, the culture can be transferred without pre-reducing the medium by using a 10% to 20% inoculum.

Notes

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and



Product Sheet

Ectothiorhodospira
***shaposhnikovii* (ATCC®)**
43036™

patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.


This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org


Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2017. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [03/13]

Please read this **FIRST**



Storage Temp.
Frozen: -80°C or colder
Freeze-Dried: 2°C to 8°C
Live Culture: See Propagation Section



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Ectothiorhodospira shaposhnikovii* (ATCC® 43036™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor