



Product Sheet

# *Desulfonispora thiosulfatigenes* (ATCC® 700533™)

Please read this **FIRST**



## Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

## Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Desulfonispora thiosulfatigenes* (ATCC® 700533™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

## Description

**Designation:** GKNTAU [DSM 11270, LK]  
**Deposited Name:** Unidentified bacterium

## Propagation

### Medium

ATCC® Medium 2188: Modified VWN medium

### Growth Conditions

**Temperature:** 30.0°C

**Atmosphere:** Under 100% N<sub>2</sub>

### Propagation Procedure

1. Sterilize the top of the Balch tube by spraying it with 70% ethanol and then flaming the top.
2. If needed, exchange the gas in the Balch tube for 100% N<sub>2</sub>.
3. If the medium is pink (see discussion about resazurin) add 2.0 ml of reducing agent (3% cysteine, stock solution) per 100 ml of medium. Let the medium sit at room temperature for 10 to 20 minutes, until the resazurin becomes colorless, before inoculating.
4. When the Balch tube is ready to inoculate, open the vial according to enclosed instructions. Take an anaerobic  
1.0 ml syringe (see discussion below) tipped with a 22 gauge needle and withdraw 0.5 ml of 2188 medium from the Balch tube and rehydrate the freeze dried pellet. Immediately place the rehydrated vial under a stream of sterile gas to maintain anaerobic conditions.
5. Using the same syringe transfer the rehydrated cell suspension back into a tube of #2188 broth. Plate 0.1 ml of the inoculated culture onto a non-selective agar medium and incubate aerobically at 30°C. Inoculate a nonselective anaerobic and aerobic broth. Transfer 0.5 ml of the rehydrated culture to a second tube of 2188 medium. Incubate the inoculated tubes at 30°C.
6. Growth should be detected in the #2188 broth within 24 to 48 hours. No growth should be detected on the aerobic plate, or in the nonselective aerobic or anaerobic broth.

### ANAEROBIC CONDITIONS:

- a. Balch tubes (available from Bellco Glass, Vineland, NJ) are specially designed for anaerobic work and use an aluminum crimp cap to hold a rubber stopper in place. Needles can easily be inserted through the stopper, and the tubes can be pressurized to 2 atm. /alternatively, serum vials may be used, or screw cap tubes with butyl rubber stoppers. In the latter case the stopper may be removed and the tube placed under a cannula system that dispenses sterile, oxygen-free gas for the addition of reducing agents or inoculation.
- b. Resazurin is a commonly used redox indicator that is pink when the redox potential is above 50 mv., and colorless when the redox potential is below 110 mv. i.e. highly reducing. Most strict anaerobes require this low redox potential for optimum growth.
- c. To obtain a fully reduced medium, it is necessary that the medium be anoxic and that a reducing agent be added. Common reducing agents are sodium sulfide, cysteine, dithiothreitol, and titanium citrate.
- d. Syringes can be made anaerobic by one of two methods. 1. Displace the dead space in the syringe with a sterile

## Notes

Cells appear as motile rods in singles and pairs with sub-terminal spores.  
Growth may be stimulated by the addition of 1 or 2 drops of dithionite (5% w/v prepared under N<sub>2</sub> and filter-sterilized) to the #2188 broth tube into which the culture is to be inoculated.

## References

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

## Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

## ATCC Warranty



Product Sheet

***Desulfonispora  
thiosulfatigenes* (ATCC®  
700533™)**

Please read this **FIRST**



**Intended Use**

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

**Citation of Strain**

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Desulfonispora thiosulfatigenes* (ATCC® 700533™)

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

**Disclaimers**

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [02/02]

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor