



Product Sheet

Aspergillus carbonarius (ATCC® 8740™)

Please read this FIRST

Storage Temp.
Frozen: -80°C or colder
Freeze-Dried: 2°C to 8°C
Live Culture: See Propagation Section

 Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Aspergillus carbonarius* (ATCC® 8740™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Strain Designation: NRRL 67 [67 (Monospore 4); 4707.878, CBS 420.64, DSM 872, IMI 41875, MUCL 30479, NRRL 1737, NRRL 605, QM 330, WB 67, Wis. (old series) 62]

Deposited Name: *Aspergillus niger* van Tieghem

Product Description: An ampoule containing viable cells (may include spores and mycelia) suspended in cryoprotectant.

Propagation

The information recommended in this section is to assist users in obtaining living culture(s) for their studies. The recommendation does not imply that the conditions or procedures provided below are optimum. Experienced researchers may initiate the growth of a culture in their own way.

ATCC® Medium 200: YM agar or YM broth

ATCC® Medium 324: Malt extract agar

ATCC® Medium 336: Potato dextrose agar (PDA)

Growth Conditions

Temperature: 24°C to 26°C

Atmosphere: Typical aerobic

Recommended Procedure

For **freeze-dry (lyophilized)** ampoules:

1. Open an ampoule according to enclosed instructions.
2. From a single test tube of **sterile distilled water** (5 to 6 mL), withdraw approximately 0.5 to 1.0 mL with a sterile pipette and apply directly to the pellet. Stir to form a suspension.
3. Aseptically transfer the suspension back into the test tube of sterile distilled water.
4. Let the test tube sit at room temperature (25°C) undisturbed for **at least 2 hours**; longer (e.g., overnight) rehydration might increase viability of some fungi.
5. Mix the suspension well. Use several drops (or make dilutions if desired) to inoculate recommended solid or liquid medium. Include a control that receives no inoculum.
6. Incubate the inoculum at the propagation conditions recommended.
7. Inspect for growth of the inoculum/strain regularly. The sign of viability is noticeable typically after 1-2 days of incubation. However, the time necessary for significant growth will vary from strain to strain.

Notes

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

DNA Sequence

18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2, complete sequence; and 28S ribosomal RNA gene, partial sequence
GGTTTCCGTAGGTGAACCTGCGGAAGGATCATTACCGAGTGC GGGTCCCTTTGGGCCCAACCTCCACCC
GTGCTATTGTACCTGTTGCTTCGGCGGGCCCGCCTTGTGCGGCCCGGGGGGATCTCTGCCCTC
GGGCCCGTGCCTCGGAGACCAACACGAACACTGTCTGAAATCGTGAAGTCTGAGTCGATTGTTT
TCAATCAGTAAAACCTTCAACAATGGATCTCTTGGTCCGGCATCGATGAAGAACGCAGCGAAATGC
GATAACTAATGTGAATTGCAGAATTCAGTGAATCATCGAGTCTTTGAACGCACATTGCGCCCTCGGTA
TTCCGGGGGGCATGCCTGTCCGAGCGTCATTGCTGCCCTCAAGCCCGGCTTGTGTGTTGGGTCGCCGTC
CCCCTGTCTGGGGGACGGGCCGAAAGGCAGCGCGGCACCGCTCCGATCCTCGAGCGTATGGGG
CTTTGTCACATGCTCTGTAGGATTGGCCGGCGCTGCCGACAACCTCAACCTTTTTTCCAGTTGACCT
CGGATCAGGTAGGATACCCGCTGAACCTAAGCATATCAATAA

D1D2 region of the 28S ribosomal RNA gene

ATATCAATAAGCGGAGGAAAAGAAACCAACCGGGATTGCCTCAGTAACGGCGAGTGAAGCGGCAAG
AGCTCAAATTTGAAAGCTGGCTCCTTCGAGTCCGCATTGTAATTTGCAGAGATGCTTTGGTGCGGC
CCCCGTCTAAGTGCCCTGGAACGGGCGGTACAGAGGGTGAGAATCCCGTCTTGGCGGGGTGTCCT
GCCCGTGTAAAGCTCCTTCGACGAGTTCGAGTTGTTGGGAATGCAGCTCTAAATGGGTGGTAAATTTCA
TCTAAAGCTAAATACTGGCCGGAGACCGATAGCGCACAAGTAGAGTGATCGAAAGATGAAAAGCAC
TTTGGAAAGAGAGTTAAACAGCACGTGAAATTTGTTGAAAGGGAAGCGCTTGAACACAGACTCGCCCG
CGGGGTTACGCCGCACTTCTGCGGGTGTACTTCCCGGTGGCGGGGCCAGCGTCCGGTTTGGCGGCCG
GTCAAAGGCCCTGGAATGTAGTGCCTCCGGGCACCTTATAGCCAGGGGTGCAATGCGGCCAGCCT
AGACCGAGGAACGCGCTTCGGCTCGGACGCTGGCATAATGTTGTAACGAC

