



Product Sheet

S. cerevisiae/E. coli marker swap vectors (ATCC® 87561™)

Please read this FIRST



Storage Temp.
Store frozen plate at -80°C



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: (ATCC® 87561™)

Shipping Information

Frozen glycerol of Escherichia coli HB101 stocks containing the plasmids arrayed on a microtiter plate

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

This is a set of 15 marker swap vectors, each containing a gene disruption conversion cassette for conversion between standard markers used for transformation selection in *Saccharomyces cerevisiae*. The vectors are useful in changing markers for gene disruptions or for changing markers on plasmids.

Maps for some of the vectors can be found on the ATCC web page or in the reference. Vectors are all in *E. coli* and arrayed on a microtiter plate. A key to the plate is provided below.

| Well | Name | ATCC | Convert | Enzymes to | Cassette | |
|------|-------|-------|-------------|------------|----------|----------------|
| | | | Function | Excise | size | Markers |
| a1 | pHL3 | 87542 | HIS3 → LEU2 | Apal+PstI | 4.3 | cmIR LEU2 |
| a2 | pHT6 | 87543 | HIS3 → TRP1 | SmaI+XhoI | 3.6 | cmIR kanR TRP1 |
| a3 | pHU10 | 87544 | HIS3 → URA3 | SmaI+XhoI | 3.2 | cmIR kanR URA3 |
| a4 | pLH7 | 87545 | LEU2 → HIS3 | XhoI+HpaI | 4.3 | cmIR kanR HIS3 |
| a5 | pLT11 | 87546 | LEU2 → TRP1 | XhoI+HpaI | 6.1 | ampR cmIR TRP1 |
| a6 | pLU12 | 87547 | LEU2 → URA3 | XhoI+HpaI | 4.1 | cmIR kanR URA3 |
| a7 | pTH4 | 87548 | TRP1 → HIS3 | XhoI+EcoRI | 3.3 | cmIR kanR HIS3 |
| a8 | pTL7 | 87549 | TRP1 → LEU2 | XhoI+SmaI | 4.4 | cmIR kanR LEU2 |
| a9 | pTU10 | 87550 | TRP1 → URA3 | XhoI+EcoRI | 3.2 | cmIR kanR URA3 |
| a10 | pUH7 | 87551 | URA3 → HIS3 | SmaI | 3.5 | ampR kanR HIS3 |
| a11 | pUL9 | 87552 | URA3 → LEU2 | SmaI | 4.6 | ampR kanR LEU2 |
| a12 | pUT11 | 87553 | URA3 → TRP1 | SmaI | 3.7 | ampR kanR TRP1 |
| b1 | M2371 | 87557 | HIS → ADE2 | PstI+SacI | 4.99 | ampR ADE2 |
| b2 | M3499 | 87558 | URA3 → ADE2 | PstI | 4.45 | ampR ADE2 |
| b3 | M2660 | 87559 | URA3 → LYS2 | HindIII | 5.57 | ampR LYS2 |

Designation: S. cerevisiae/E. coli marker swap vectors

Notes

To convert the host phenotype from the existing yeast auxotrophic marker to a new marker (eg. HIS3 → LEU2), transform with the restriction enzyme digested vector (eg. Apal+PstI digested pHL3) and select for the appropriate phenotype (for pHL3, Leu+). Some combinations of marker swap plasmids and target locus may result in relatively high reversion rates. In most, but not all cases, the frequencies of successful convertants are greater than 30%. When swapping markers on an episomal plasmid, appropriate phenotype may result from loss of the plasmid unless a second selectable or scorable marker is used to ensure plasmid maintenance. Refer to reference for more information.
- Yeast 13: 647-653, 1997

Vector Information

ATCC® Catalog No. 87542
Designation: pHL3 Other name(s): HIS3 → LEU2 converter Excise insert: Apal+PstI
Size of construct (kb): 7.7
Markers: cmIR, LEU2
Growth temperature: 37°C
Growth medium: LB medium + 25 µg/mL chloramphenicol

NOTES:
Restriction digests of the clone give the following sizes (kb): BgIII--4.4, 3.1; EcoRI--7.6; HindIII--4.1, 3.5.
- ATCC staff
To convert the host phenotype from HIS3 to LEU2, transform with the Apal+PstI digested vector and select for Leu+ transformants. Vector was constructed by replacing an internal BgIII fragment of HIS3 with a 2.7 kb BgIII fragment containing the LEU2 coding sequence. LEU2 and HIS3 are in the same orientation.
- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87543
Designation: pHT6
Other name(s): HIS3 → TRP1 converter
Excise insert: SmaI+XhoI
Size of construct (kb): 7.0
Markers: cmIR, kanR, TRP1
Growth temperature: 37°C



Product Sheet

***S. cerevisiae*/E. coli marker swap vectors (ATCC® 87561™)**

Please read this FIRST

| | |
|---|---|
|  | Storage Temp. Store frozen plate at -80°C |
|  | Biosafety Level 1 |

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: (ATCC® 87561™)

Shipping Information

Frozen glycerol of *Escherichia coli* HB101 stocks containing the plasmids arrayed on a microtiter plate

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Growth medium: LB + kanamycin (50 µg/mL) + chloramphenicol (25 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): HindIII/PstI--5.5, 1.0, 0.6; HindIII--7.2; HindIII/SalI--5.1, 2.2.

- ATCC staff

To convert the host phenotype from HIS3 to TRP1, transform with the SmaI+XhoI digested vector and select for Trp+ transformants. Vector was constructed by replacing an internal HindIII fragment of HIS3 with a SmaI fragment containing the TRP1 and kanR coding sequences. TRP1 and HIS3 are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87544

Designation: pHU10

Other name(s): HIS3 → URA3 converter

Excise insert: SmaI+XhoI

Size of construct (kb): 6.6

Markers: cmlR, kanR, URA3

Growth temperature: 37°C

Growth medium: LB + kanamycin (50 µg/mL) + chloramphenicol (25 µg/mL)

NOTES: Restriction digests of the clone give the following sizes (kb): HindIII--6.5; HindIII/PstI--5.0, 1.1, 0.5.

- ATCC staff

To convert the host phenotype from HIS3 to URA3, transform with the SmaI+XhoI digested vector and select for Ura+ transformants. Vector was constructed by replacing an internal HindIII fragment of HIS3 with a SmaI fragment containing the URA3 and kanR coding sequences. URA3 and HIS3 are in the opposite orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87545

Designation: pLH7

Other name(s): LEU2 → HIS3 converter

Excise insert: XhoI+HpaI

Size of construct (kb): 8.7

Markers: cmlR, kanR, HIS3

Growth temperature: 37°C

Growth medium: LB + kanamycin (50 µg/mL) + chloramphenicol (25 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): BamHI/EcoRI--6.6, 2.0; XhoI--8.7.

- ATCC staff

To convert the host phenotype from LEU2 to HIS3, transform with the XhoI+HpaI digested vector and select for His+ transformants. Vector was constructed by replacing an internal EcoRV fragment of LEU2 with a SmaI fragment containing the HIS3 and kanR coding sequences. HIS3 and LEU2 are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87546

Designation: pLT11

Other name(s): LEU2 → TRP1 converter

Excise insert: XhoI+HpaI

Size of construct (kb): 10.5

Markers: cmlR, ampR, TRP1

Growth temperature: 37°C

Growth medium: LB + ampicillin (50 µg/mL) + chloramphenicol (20 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): BamHI--8.4, 2.1; BamHI/XhoI--4.9, 3.3, 2.2.

- ATCC staff

To convert the host phenotype from LEU2 to TRP1, transform with the XhoI+HpaI digested vector and select for Trp+ transformants. Vector was constructed by replacing an internal EcoRI-EcoRV fragment of LEU2 with an XhoI-EcoRI fragment containing the TRP1 and ampR coding sequences.

- Yeast 13: 647-653, 1997



Product Sheet

S. cerevisiae/*E. coli* marker swap vectors (ATCC® 87561™)

Please read this FIRST

| | |
|---|---|
|  | Storage Temp. Store frozen plate at -80°C |
|  | Biosafety Level 1 |

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: (ATCC® 87561™)

Shipping Information

Frozen glycerol of *Escherichia coli* HB101 stocks containing the plasmids arrayed on a microtiter plate

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

ATCC® Catalog No. 87547

Designation: pLU12

Other name(s): LEU2 → URA3 converter

Excise insert: XhoI/HpaI

Size of construct (kb): 8.5

Markers: cmlR, kanR, URA3

Growth temperature: 37°C

Growth medium: LB + kanamycin (50 µg/mL) + chloramphenicol (25 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): BamHI--8.8; BamHI/XhoI--5.5, 3.2; EcoRI/HindIII--7.5, 1.2; HpaI/XhoI--4.6, 4.2.

- ATCC staff

To convert the host phenotype from LEU2 to URA3, transform with the XhoI/HpaI digested vector and select for Ura+ transformants. Vector was constructed by replacing an internal EcoRV fragment of LEU2 with a SmaI fragment containing the URA3 and kanR coding sequences. URA3 and LEU2 are in the opposite orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87548

Designation: pTH4

Other name(s): TRP1 → HIS3

Excise insert: XhoI+EcoRI

Size of construct (kb): 6.7

Markers: cmlR, kanR, HIS3

Growth temperature: 37°C

Growth medium: LB + kanamycin (50 µg/mL) plus chloramphenicol (25 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): EcoRI--6.7; HindIII--4.8, 1.9; EcoRI/XhoI--3.5, 3.3.

- ATCC staff

To convert the host phenotype from TRP1 to HIS3, transform with the XhoI+EcoRI digested vector and select for His+ transformants. Vector was constructed by replacing an internal EcoRV fragment of TRP1 with a SmaI fragment containing the HIS3 and kanR coding sequences. HIS3 and TRP1 are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87549

Designation: pTL7

Other name(s): TRP1 → LEU2 converter

Excise insert: XhoI+SmaI

Size of construct (kb): 7.8

Markers: cmlR, kanR, LEU2

Growth temperature: 37°C

Medium composition: LB + kanamycin (50 µg/mL) + chloramphenicol (25 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): EcoRI--6.2, 1.7; HindIII--7.8; SmaI/XhoI--4.4, 3.3.

- ATCC staff

To convert the host phenotype from TRP1 to LEU2, transform with the XhoI+SmaI digested vector and select for Leu+ transformants. Vector was constructed by replacing an internal EcoRV fragment of TRP1 with a SmaI fragment containing the LEU2 and kanR coding sequences. LEU2 and TRP1 are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87550

Designation: pTU10



Product Sheet

***S. cerevisiae*/E. coli marker swap vectors (ATCC® 87561™)**

Please read this FIRST

| | |
|---|---|
|  | Storage Temp. Store frozen plate at -80°C |
|  | Biosafety Level 1 |

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: (ATCC® 87561™)

Shipping Information

Frozen glycerol of *Escherichia coli* HB101 stocks containing the plasmids arrayed on a microtiter plate

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Other name(s): TRP1→ URA3 converter

Excise insert: XhoI+EcoRI

Size of construct (kb): 6.6

Markers: *cmIR*, *kanR*, *URA3*

Growth temperature: 37°C

Medium composition: LB + kanamycin (50 µg/mL) plus chloramphenicol (25 µg/mL)

NOTES:

Restriction digests of the clone give the following sizes (kb): EcoRI--6.6; HindIII--5.0, 1.6; EcoRI/XhoI--3.4 3.2.

- ATCC staff

To convert the host phenotype from TRP1 to URA3, transform with the XhoI+EcoRI digested vector and select for Ura⁺ transformants. Vector was constructed by replacing an internal EcoRV fragment of TRP1 with a SmaI fragment containing the URA3 and kanR coding sequences. URA3 and TRP1 are in the opposite orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87551

Designation: pUH7

Other name(s): URA3 → HIS3 converter

Excise insert: SmaI

Size of construct (kb): 6.5

Markers: *ampR*, *kanR*, *HIS3*

Growth temperature: 37°C

Medium composition: LB medium + 50 µg/mL ampicillin and 20 µg/mL kanamycin

NOTES:

Restriction digests of the clone give the following sizes (kb): EcoRI--6.4; SmaI--3.5, 2.8; XbaI--3.6, 2.7.

- ATCC staff

To convert the host phenotype from URA3 to HIS3, transform with the SmaI digested vector and select for His⁺ transformants. Vector was constructed by replacing an internal *StuI* fragment of URA3 with a SmaI fragment containing the HIS3 and kanR coding sequences. HIS3 and URA3 are probably in the opposite orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87552

Designation: pUL9

Other name(s): URA3 → LEU2 converter

Excise insert: SmaI

Size of construct (kb): 7.45

Markers: *ampR*, *kanR*, *LEU2*

Growth temperature: 37°C

Growth medium: LB medium + 50 µg/mL ampicillin and 20 µg/mL kanamycin

NOTES:

Restriction digests of the clone give the following sizes (kb):

Expected SmaI digest fragments of 2.74 kb and 4.71 kb

Expected EcoRI digest fragments of 2.95 kb and 4.5 kb

- ATCC staff

To convert the host phenotype from URA3 to LEU2, transform with the SmaI digested vector and select for Leu⁺ transformants. Vector was constructed by replacing an internal *StuI* fragment of URA3 with a SmaI fragment containing the LEU2 and kanR coding sequences. LEU2 and URA3 are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87553

Designation: pUT11

Other name(s): URA3 → TRP1 converter

Excise insert: SmaI

Size of construct (kb): 6.7

Markers: *ampR*, *kanR*, *TRP1*

Growth temperature: 37°C

Growth medium: LB medium + 50 µg/mL ampicillin and 20 µg/mL kanamycin



Product Sheet

***S. cerevisiae*/E. coli marker swap vectors (ATCC® 87561™)**

Please read this FIRST

| | |
|---|---|
|  | Storage Temp. Store frozen plate at -80°C |
|  | Biosafety Level 1 |

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: (ATCC® 87561™)

Shipping Information

Frozen glycerol of *Escherichia coli* HB101 stocks containing the plasmids arrayed on a microtiter plate

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

NOTES:

Restriction digests of the clone give the following sizes (kb): *HindIII*--4.6, 2.0; *SmaI*--3.8, 2.7.

- ATCC staff

To convert the host phenotype from *URA3* to *TRP1*, transform with the *SmaI* digested vector and select for *Trp+* transformants. Vector was constructed by replacing an internal *StuI* fragment of *URA3* with a *SmaI* fragment containing the *TRP1* and *kanR* coding sequences. *TRP1* and *URA3* are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87557

Designation: M2371

Other name(s): *HIS3* → *ADE2* converter

Excise insert: *PstI*+ *SacI*

Size of construct (kb): 7.734

Markers: *ampR* *ADE2*

Growth temperature: 37°C

Growth medium: LB Medium + 50 µg/mL ampicillin

NOTES:

Restriction digests of the clone give the following sizes (kb): *BamHI*--3.7, 2.7, 0.7, 0.6; *PvuI*--6.8, 0.9;

HindIII--3.9, 1.6, 1.2, 1.0.

- ATCC staff

To convert the host phenotype from *HIS3* to *ADE2*, transform with the *PstI*+*SacI* digested vector (4.99 kb) and select for *Ade+* transformants. Vector was constructed by replacing an internal *MscI*-*NsiI* fragment of *HIS3* with a 3.7 kb *NotI* (blunt)-*PstI* fragment containing the *ADE2* coding sequence. *ADE2* and *HIS3* are in the opposite orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87558

Designation: M3499

Other name(s): *URA3* → *ADE2* converter

Excise insert: *PstI*

Size of construct (kb): 7.426

Markers: *ampR* *ADE2*

Growth temperature: 37°C

Growth medium: LB Medium + 50 µg/mL ampicillin

NOTES:

Restriction digests of the clone give the following sizes (kb): *HindIII*--2.5, 1.3, 1.1 (doublet), 1.0; *BamHI*--6.6, 0.5; *PvuI*--6.2, 0.8.

- ATCC staff

To convert the host phenotype from *URA3* to *ADE2*, transform with the *PstI* digested vector (4.45 kb) and select for *Ade+* transformants. Vector was constructed by replacing an internal *StuI*-*EcoRV* fragment of *URA3* with a 3.7 kb *BamHI* fragment containing the *ADE2* coding sequence. *ADE2* and *URA3* are in the same orientation.

- Yeast 13: 647-653, 1997

ATCC® Catalog No. 87559

Designation: M2660

Other name(s): *URA3* → *LYS2*

Excise insert: *HindIII*

Size of construct (kb): 8.308

Markers: *ampR* *LYS2*

Growth temperature: 37°C

Growth medium: LB Medium + 50 µg/mL ampicillin

NOTES:

Restriction digests of the clone give the following sizes (kb): *BglII*--4.3, 4.1; *HindIII*--5.6, 2.8; *HpaI*--8.5; *PvuI*--5.1, 2.4, 0.9.

- ATCC staff

To inactivate *URA3*, transform with *HindIII*-digested plasmid (5.57 kb) and select for *Lys+*. Constructed by



Product Sheet

***S. cerevisiae*/E. coli marker swap vectors (ATCC® 87561™)**

Please read this FIRST



Storage Temp.
Store frozen plate at -80°C



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: (ATCC® 87561™)

Shipping Information

Frozen glycerol of *Escherichia coli* HB101 stocks containing the plasmids arrayed on a microtiter plate

replacing an internal EcoRV-PstI fragment of URA3 with a 4.6 kb PvuII-PstI fragment containing LYS2. URA3 and LYS2 are in the same orientation.

- Yeast 13: 647-653, 1997

Propagation

Transfer a loopful to a test tube containing 5 mL LB+ antibiotic (either 50 µg/mL of ampicillin 10 µg/mL chloramphenicol or 25 µg/mL chloramphenicol and 50 µg/mL kanamycin check specific markers). A loopful of culture can also be streaked on an LB + amp or chl agar plate. Incubate cultures at 37°C. Isolate DNA using standard plasmid preparation procedures.

Growth Conditions

Temperature: 37°C

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the Biosafety in Microbiological and Biomedical Laboratories from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2016. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [01/11]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor