



## Product Sheet

# 293T/17 SF [HEK 293T/17 SF] (ATCC® ACS-4500™)

### Please read this FIRST



Storage Temp.  
**liquid nitrogen**  
vapor phase

---



Biosafety Level  
**2**

### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Complete Growth Medium

The base medium for this cell line is IS 293-V (Irvine Scientific No. 91107). To make complete medium, add:

- 200 mM L-Alanyl-L-Glutamine, 200 mM Solution in 0.85% NaCl (ATCC® No. 30-2115™) to a final concentration of 8 mM
- 10% Poloxamer 188 (Sigma P5556) to a final concentration of 0.1%
- 10uL/mL of ITS Solution (Insulin, Transferrin, Selenious Acid, Corning #25-800-CR)  
This medium is formulated for use with a 5-8% CO<sub>2</sub> air atmosphere.

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: 293T/17 SF [HEK 293T/17 SF] (ATCC® ACS-4500™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

### Description

**Organism:** *Homo sapiens*, human

**Tissue:** Kidney

**Morphology:** Lymphocyte-like; single cells to small aggregates

**Growth Properties:** suspension

### Batch-Specific Information

Refer to the Certificate of Analysis for batch-specific test results.

### SAFETY PRECAUTION

ATCC highly recommends that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

### Unpacking & Storage Instructions

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

### Handling Procedure for Frozen Cells

To ensure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If, upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at ?80°C. Storage at 80°C will result in loss of viability.

1. Rapidly thaw cells by placing the cryovial in a 37°C water bath, swirling gently. Remove the cryovial from the water bath when only a few ice crystals are remaining.
2. Sterilize the cryovial by rinsing with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. Using a 1-mL or 5-mL pipette, transfer thawed cells drop-wise into 9 mL pre-warmed complete growth medium in a 15-mL centrifuge tube. Gently pipette the cells up and down several times to mix thoroughly.
4. Centrifuge the cell suspension at 170 × g for 5 minutes.
5. Carefully aspirate the supernatant and discard, leaving the cell pellet.
6. Gently resuspend the cell pellet in fresh pre-warmed complete growth medium, and transfer cell suspension into a filtered cap/non-baffled shaker flask. Cells should be seeded at a density of 5 × 10<sup>5</sup> cells/mL.
7. Place the flask in a 37°C shaking incubator (125 to 130 rpm) with 5-8% CO<sub>2</sub>. **Note:** Viability after initial thaw is generally lower; however, after 2-3 passages, the cells are fully recovered and reach optimal viability.

### Subculturing Procedure

Subculture cells at log phase (when cells are ready for passaging, i.e., every 2-3 days, and are approximately 2 × 10<sup>6</sup> cells/mL). Pre-warm fresh growth medium prior to use. Swirl the flask gently to evenly distribute cells in medium. Remove a small volume of cells from the flask and perform cell count.

1. Seed at 5×10<sup>5</sup> cells/mL for a 2 day subculture and 4×10<sup>5</sup> cells/mL for a 3 day subculture (weekend)
2. To maintain high cell viability, prior to seeding, centrifuge cells for 5min at 170x g
3. Discard spent media and re-suspend cell pellet in pre-warmed fresh complete growth media
4. Pipette cells gently to break aggregates

Note: Slight aggregates may be observed, but they are easily dispersed with minimal pipetting and do not impact the performance of the cell line. Alternately, appropriate amount of fresh media maybe added directly into the flask to adjust cell seeding density. However, cell viability might be slightly compromised and decreased by 5%.

### Cryopreservation Medium

Prepare 2X freezing medium: Complete Growth Medium supplemented with 15% (v/v) DMSO.


### Comments



## Product Sheet


# 293T/17 SF [HEK 293T/17 SF] (ATCC® ACS-4500™)

### Please read this FIRST



Storage Temp.  
**liquid nitrogen**  
vapor phase

---



Biosafety Level  
**2**

### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Complete Growth Medium

The base medium for this cell line is IS 293-V (Irvine Scientific No. 91107). To make complete medium, add:

- 200 mM L-Alanyl-L-Glutamine, 200 mM Solution in 0.85% NaCl (ATCC® No. 30-2115™) to a final concentration of 8 mM
- 10% Poloxamer 188 (Sigma P5556) to a final concentration of 0.1%
- 10uL/mL of ITS Solution (Insulin, Transferrin, Selenious Acid, Corning #25-800-CR)  
This medium is formulated for use with a 5-8% CO<sub>2</sub> air atmosphere.

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: 293T/17 SF [HEK 293T/17 SF] (ATCC® ACS-4500™)

The cells constitutively express the temperature-sensitive SV40 T antigen that allows for episomal replication of transfected plasmids containing the SV40 origin of replication. This feature increases protein expression levels by permitting more plasmid copies to persist in the transiently transfected cells. Expression vectors containing the human cytomegalovirus (CMV) promoter have been shown to achieve high levels of protein expression in 293T/17 cell line.

Transient transfections can be performed at small and large scale. High transfection efficiencies and protein yields have been demonstrated in this cell line. ATCC recommends passaging thawed cells at least twice prior to transfection to ensure optimal viability. Prior to transfection (24 hours), seed cells at a density of  $8 \times 10^5$  cells/mL.

### References

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

### Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

### ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

### Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate. This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials. Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).  
© ATCC 2017. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [05/25]

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor