



Product Sheet

Mixed culture (ATCC® BAA-845™)

Please read this FIRST



Storage Temp.
Frozen: -80°C or colder
Freeze-Dried: 2°C to 8°C
Live Culture: See Propagation Section



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Mixed culture* (ATCC® BAA-845™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Designation: 5.2 s.c.1

Deposited Name: Unidentified bacterium

Product Description: Mixed culture of *Chroogloeocystis* sp. and *Silanimonas lenta*

Propagation

Medium

ATCC® Medium 2453: DH medium

Growth Conditions

Temperature: 45°C to 55°C

Atmosphere: Aerobic, under light intensity of 500 to 1,000 lux

Propagation Procedure

1. Withdraw 0.6 mL from the base of a broth culture where cells are concentrated, or harvest cells from a slant culture with 0.6 mL of #2453 broth.
2. Using this aliquot, inoculate one broth and one slant tube with 0.2 and 0.4 mL respectively.
3. Incubate tubes at 55°C under 2000-3000 LUX light.

Notes

Good growth, indicated by increased pigmentation in the broth or on the slant, should occur after one to two weeks of incubation. Examine cells microscopically to assure that they are intact and healthy. At this time additional test tubes or flasks can be inoculated. A 10% inoculum is recommended (i.e. 10.0 mL of culture to 100.0 mL fresh medium).

To minimize change in a culture, it is recommended that a frozen seed stock be established from early passage cells. This may be accomplished by propagating the strain under ideal conditions, utilizing recommended medium, temperature and light. Prepare a concentrated cell suspension, after good growth is achieved. If grown in broth, pellet the cells by centrifugation. Decant the supernatant and resuspend the pellet in fresh #2453 broth using 1/10 or less of the original volume. For slant cultures, wash cells off the agar surface with a minimal amount of #2453 broth so that a concentrated cell suspension is attained. Add 50% DMSO solution to the concentrated cell suspension so that the final concentration of DMSO in the suspension is 5%. Dispense small aliquots (0.5 to 1 mL) of the suspension into small sterile vials. Store the vials at -50°C or below.

When needed, remove vials from storage, thaw contents in a 37°C water bath and inoculate into recommended medium. A minimum of 0.2 mL of the thawed stock should be used to inoculate 5 mL of broth or 1 agar slant.

Additional information on this culture is available on the ATCC® web site at www.atcc.org.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.


Disclaimers




Product Sheet

Mixed culture (ATCC® BAA-845™)

Please read this FIRST



Storage Temp.
**Frozen: -80°C or
colder**
**Freeze-Dried: 2°C
to 8°C**
**Live Culture: See
Propagation
Section**



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Mixed culture* (ATCC® BAA-845™)

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2017. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [08/24]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor