




Product Sheet

Desulfovibrio gracilis (ATCC® BAA-904™)

Please read this **FIRST**



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Desulfovibrio gracilis* (ATCC® BAA-904™)

Description

Designation: SRL6146 [DSM 16080]

Propagation

Medium

ATCC® Medium 2465: *Desulfovibrio gracilis* medium

Growth Conditions

Temperature: 37.0°C

Atmosphere: Anaerobic

Propagation Procedure

1. Open the vial according to enclosed instructions.
2. Perform all steps under anaerobic conditions.
3. Aseptically transfer 0.5 ml of #2465 broth to the vial and rehydrate the pellet. Transfer the suspension back into the broth tube. Inoculate a plate of a non-selective medium such as Tryptic Soy, Nutrient, or blood agar with 0.1 ml of the cell suspension.
4. Seal the tube with a rubber stopper and incubate anaerobically at 37°C. Incubate the plate(s) aerobically as a purity check.
5. After one or two days, growth should be evident as indicated by turbidity through out the broth. Once growth has been established the culture should be transferred to fresh broth every 24 to 48 hours.
6. This culture is very sensitive to oxygen when initially rehydrated, therefore steps should be taken to avoid exposure to oxygen. When the culture exhibits good growth it will remain viable for up to 1 week if stored at 4°C under anaerobic condition.

ANAEROBIC CONDITIONS:

- Tubes of media are placed under a gassing cannula system hooked to a source of oxygen free gas.
- All transfers are performed while the test tubes are on the cannula system with a gentle stream of oxygen free gas flowing through the system.
- As the test tubes are removed from the cannula system each is sealed with butyl rubber stopper thus maintaining the anaerobic headspace.
- 100% nitrogen or 80% nitrogen-20% carbon dioxide gas mixture is typically employed as the oxygen free gas source.

Notes

When examined microscopically, the cells appear as single (some pairs) long straight rods that are motile. Always use freshly prepared anaerobic medium. If there is any question about the anaerobic condition of the medium, the medium can be reduced with the addition of 1.5% cysteine (2.0 ml per 100 ml of medium). Other commonly used reducing agents are sodium sulfide, cysteine, dithiothreitol, and titanium citrate. Cysteine is the reducing agent of choice since it does not cause the ferrous ammonium sulfate to precipitate.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor



Product Sheet

Desulfovibrio gracilis **(ATCC® BAA-904™)**

Please read this FIRST



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Desulfovibrio gracilis* (ATCC® BAA-904™)

longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2012. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [12/04]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor