



Product Sheet

MeT-5A (ATCC® CRL-9444™)

Please read this FIRST

Storage Temp.
**liquid nitrogen
vapor phase**

Biosafety Level
2

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Complete Growth Medium

The base medium for this cell line is Medium 199 containing 1.5 g/L sodium bicarbonate.

To make the complete growth medium, add the following components to the base medium:

10% fetal bovine serum (final conc.)

3.3 nM epidermal growth factor (EGF) (final conc.) (do not filter).

400 nM hydrocortisone (final conc.)

870 nM zinc-free bovine insulin (final conc.)

20 mM HEPES (final conc.)

The trace elements at the following final concentrations:

- H_2SeO_3 0.3869 mg/L (Selenious acid)
- $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ 0.0198 mg/L (Manganese chloride)
- $\text{Na}_2\text{SiO}_3 \cdot 9\text{H}_2\text{O}$ 14.2100 mg/L (Sodium silicate)
- $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ 0.1236 mg/L (Ammonium molybdate)
- NH_4VO_3 0.0585 mg/L (Ammonium vanadate)
- $\text{NiSO}_4 \cdot 6\text{H}_2\text{O}$ 0.0131 mg/L (Nickel sulfate)
- $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ 0.0113 mg/L (Tin Chloride)

See Methods in Cell Biology, Vol. 21B, pg. 200, 1980

American Type Culture Collection

PO Box 1549

This medium is formulated for use with a 5% CO_2 in air

atmosphere.

ATCC tested fetal bovine serum is available as ATCC

Catalog No. 30-2020 (500ml) and ATCC Catalog No.

59-2729 (100ml).

Email: Tech@atcc.org

Citation of Strain Distributor

Description

Organism: *Homo sapiens*, human

Tissue: mesothelium

Disease: pleural fluids obtained from non-cancerous individuals.

Cell Type: epithelial virus transformed

Morphology: epithelial

Growth Properties: adherent

DNA Profile:

Amelogenin: X,Y

CSF1PO: 10,12

D13S317: 11,13

D16S539: 12

D5S818: 12

D7S820: 10

THO1: 6,9,3

TPOX: 8

vWA: 15,18

Batch-Specific Information

Refer to the Certificate of Analysis for batch-specific test results.

SAFETY PRECAUTION

ATCC highly recommends that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

Unpacking & Storage Instructions

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C , preferably in liquid nitrogen vapor, until ready for use.

Handling Procedure for Frozen Cells

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. It is recommended that the cryoprotective agent be removed immediately. Centrifuge the cell suspension at approximately $125 \times g$ for 5 to 10 minutes. Discard the supernatant and resuspend the cell pellet in an appropriate amount of fresh growth medium.
4. Transfer the vial contents to an appropriate size vessel. It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).
5. Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Handling Procedure for Flask Cultures

The flask was seeded with cells (see specific batch information) grown and completely filled with medium at ATCC to prevent loss of cells during shipping.

1. Upon receipt visually examine the culture for macroscopic evidence of any microbial contamination. Using an inverted microscope (preferably equipped with phase-contrast optics), carefully check for any evidence of microbial contamination. Also check to determine if the majority of cells are still attached to the bottom of the flask; during shipping the cultures are sometimes handled roughly and



Product Sheet

MeT-5A (ATCC® CRL-9444™)

Please read this FIRST

Storage Temp.

liquid nitrogen

vapor phase

Biosafety Level

2

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Complete Growth Medium

The base medium for this cell line is Medium 199 containing 1.5 g/L sodium bicarbonate.

To make the complete growth medium, add the following components to the base medium:

- 10% fetal bovine serum (final conc.)
- 3.3 nM epidermal growth factor (EGF) (final conc.) (do not filter).
- 400 nM hydrocortisone (final conc.)
- 870 nM zinc-free bovine insulin (final conc.)
- 20 mM HEPES (final conc.)

The trace elements at the following final concentrations:

- H₂SeO₃ 0.3869 mg/L (Selenious acid)
- MnCl₂•4H₂O 0.0198 mg/L (Manganese chloride)
- Na₂SiO₃•9H₂O 14.2100 mg/L (Sodium silicate)
- (NH₄)₆Mo₇O₂₄•4H₂O 0.1236 mg/L (Ammonium molybdate)
- NH₄VO₃ 0.0585 mg/L (Ammonium vanadate)
- NiSO₄•6H₂O 0.0131 mg/L (Nickel sulfate)
- SnCl₂•2H₂O 0.0113 mg/L (Tin Chloride)

See Methods in Cell Biology, Vol. 21B, pg. 200, 1980 American Type Culture Collection

PO Box 1549
This medium is formulated for use with a 5% CO₂ in air atmosphere.
Manassas, VA 20108 USA
www.atcc.org

ATCC tested fetal bovine serum is available as ATCC Catalog No. 30-2020 (500ml) and ATCC Catalog No. 50-2020 (100ml).

50-2020 (100ml)
Email: Tech@atcc.org

Citation of Strain

- many of the cells often detach and become suspended in the culture medium (but are still viable).
- 2. **If the cells are still attached**, aseptically remove all but 5 to 10 mL of the shipping medium. The shipping medium can be saved for reuse. Incubate the cells at 37°C in a 5% CO₂ in air atmosphere until they are ready to be subcultured.
- 3. **If the cells are not attached**, aseptically remove the entire contents of the flask and centrifuge at 125 x g for 5 to 10 minutes. Remove shipping medium and save. Resuspend the pelleted cells in 10 mL of this medium and add to 25 cm² flask. Incubate at 37°C in a 5% CO₂ in air atmosphere until cells are ready to be subcultured.



Subculturing Procedure

Volumes are given for a 75 cm² flask. Increase or decrease the amount of dissociation medium needed proportionally for culture vessels of other sizes.

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin- 0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting.
5. To remove trypsin-EDTA solution, transfer cell suspension to centrifuge tube and spin at approximately 125 x g for 5 to 10 minutes. Discard supernatant and resuspend cells in fresh growth medium. Add appropriate aliquots of cell suspension to new culture vessels.
6. Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:2 to 1:4 is recommended

Medium Renewal: Every 2 to 3 days



Cryopreservation Medium

Complete culture medium described above supplemented with 5% (v/v) DMSO. Cell culture tested DMSO is available as ATCC Catalog No. 4-X.



Comments

The cells stain positively for vimentin, keratins and SV40 T antigen.



References

References and other information relating to this product are available online at www.atcc.org.



Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been



Product Sheet

MeT-5A (ATCC® CRL-9444™)

Please read this FIRST



Storage Temp.
liquid nitrogen
vapor phase



Biosafety Level
2

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Complete Growth Medium

The base medium for this cell line is Medium 199 containing 1.5 g/L sodium bicarbonate.

To make the complete growth medium, add the following components to the base medium:

10% fetal bovine serum (final conc.)

3.3 nM epidermal growth factor (EGF) (final conc.) (do not filter).

400 nM hydrocortisone (final conc.)

870 nM zinc-free bovine insulin (final conc.)

20 mM HEPES (final conc.)

The trace elements at the following final concentrations:

- H_2SeO_3 0.3869 mg/L (Selenious acid)
- $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ 0.0198 mg/L (Manganese chloride)
- $\text{Na}_2\text{SiO}_3 \cdot 9\text{H}_2\text{O}$ 14.2100 mg/L (Sodium silicate)
- $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ 0.1236 mg/L (Ammonium molybdate)
- NH_4VO_3 0.0585 mg/L (Ammonium vanadate)
- $\text{NiSO}_4 \cdot 6\text{H}_2\text{O}$ 0.0131 mg/L (Nickel sulfate)
- $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ 0.0113 mg/L (Tin Chloride)

See Methods in Cell Biology, Vol. 21B, pg. 200, 1980
American Type Culture Collection

PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

ATCC tested fetal bovine serum is available as ATCC
Catalog No. 30-2020 (500ml) and ATCC Catalog No.

59-2021 (500ml)
Email: Tech@atcc.org

Citation of Strain Distributor

confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2017. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [04/03]