



## Product Sheet

# ILB1-H6 [ILB1-H6A1B] (ATCC® HB-10219™)

### Please read this FIRST



### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Complete Growth Medium

The base medium for this cell line is HB101 medium (from Irvine Scientific). To make the complete growth medium, add fetal bovine serum to a final concentration of 7%. Alternately, cells may be grown on ATCC Hybri-Care Medium Catalog No. 46-X as the base medium. Hybri-Care Medium is supplied as a powder and should be reconstituted in 1 L cell-culture-grade water. It must be supplemented with 1.5 g/L sodium bicarbonate.

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: ILB1-H6 [ILB1-H6A1B] (ATCC® HB-10219™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

### Description

**Organism:** *Mus musculus* (B cell); *Mus musculus* (myeloma), mouse (B cell); mouse (myeloma)

**Isotype:** IgG1; kappa light chain

**Cell Type:** hybridoma: B lymphocyte

**Morphology:** lymphoblast

**Growth Properties:** suspension

### Batch-Specific Information

Refer to the Certificate of Analysis for batch-specific test results.

### SAFETY PRECAUTION

ATCC highly recommends that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

### Unpacking & Storage Instructions

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

### Handling Procedure for Frozen Cells

#### Part A. FROZEN CELLS

Vol./Ampule: 1.0 mL.

Recommended Handling Upon Receipt: Initiate culture as soon as possible upon receipt. Thaw by rapid agitation in 37°C water bath. See instructions on back.

Dilute ampule contents with culture medium (see batch data above). Add fresh medium (depending on cell density) every 2-3 days.

### Handling Procedure for Flask Cultures

#### Part B. FLASK CULTURES

Recommended Handling Upon Receipt:

Suspension Cultures: The culture flask was seeded, see batch data above, and completely filled with medium to prevent loss of cells in transit. Upon receipt incubate the flask in an upright position for several hours to return the flask contents to 37°C. After the temperature has equilibrated, aseptically remove the entire contents of the flask and centrifuge at 300 x g for 15 minutes. Resuspend the cell pellet in 10-12 mL of the shipping medium. From this suspension remove a sample for a cell count and viability so that the cell density of the suspension can be adjusted to 2-5 x 10<sup>5</sup> viable cells/mL. If the suspension needs to be diluted use the shipping medium. Incubate the culture in a flat position at 37°C in a 5% CO<sub>2</sub> in air atmosphere. Maintain the cell density of the culture as suggested under the subculture procedure described above.

### Subculturing Procedure

**Medium Renewal:** Every 2 to 3 days

Cultures can be maintained by addition or replacement of fresh medium. Start cultures at 2 X 10<sup>5</sup> cells/ml and maintain between 1 X 10<sup>5</sup> and 1 X 10<sup>6</sup> cells/ml.

### Comments

Animals were immunized with purified recombinant human IL-1 beta (17500 dalton form).

Spleen cells were fused with P3X63Ag8.653 myeloma cells.

The antibody reacts specifically with IL-1 beta and does not react with IL-1 alpha or with acidic or basic fibroblast growth factor.

It recognizes an epitope distinct from that bound by ILB1-H21 and ILB1-H67 (ATCC HB-10220 and ATCC HB-



Product Sheet

## ILB1-H6 [ILB1-H6A1B] (ATCC® HB-10219™)

Please read this **FIRST**



### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Complete Growth Medium

The base medium for this cell line is HB101 medium (from Irvine Scientific). To make the complete growth medium, add fetal bovine serum to a final concentration of 7%. Alternately, cells may be grown on ATCC Hybri-Care Medium Catalog No. 46-X as the base medium. Hybri-Care Medium is supplied as a powder and should be reconstituted in 1 L cell-culture-grade water. It must be supplemented with 1.5 g/L sodium bicarbonate.

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: ILB1-H6 [ILB1-H6A1B] (ATCC® HB-10219™)

10222) and that may overlap the epitope bound by ILB1-H34 (ATCC HB-10221). The affinity constant of ILB1-H6 was measured to be  $0.78 \times 10^{10}$  liter/mol.

### References

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

### Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

### ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

### Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate. This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials. Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).  
© ATCC 2015. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [10/19]

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor