


Product Sheet

Prostate Epithelial Cell Basal Medium (ATCC[®] PCS-440-030[™])

Please read this FIRST



American Type Culture Collection PO Box 1549 Manassas, VA 20108 USA www.atcc.org

800.638.6597 or 703.365.2700 Fax: 703.365.2750 Email: <u>Tech@atcc.org</u>

Or contact your local distributor

Product Description:

Prostate Epithelial Cell Basal Medium is a sterile, phenol red-free, liquid tissue culture medium intended for use as one component in a complete ATCC [®] Primary Cell Solutions™ system. This serum-free system is designed to support prostate epithelial cells derived from normal human prostate. Prostate Epithelial Cell Basal Medium contains essential and non-essential amino acids, vitamins, other organic compounds, trace minerals and inorganic salts. To support the proliferation and plating efficiency of various types of prostate epithelial cells, Prostate Cell Basal Medium must be supplemented with the appropriate cell-specific growth kit. When using this complete media system, the growth of prostate epithelial cells is supported without the use of feeder layers, extracellular matrix proteins or other substrates.

- For prostate epithelial cells derived from prostate tissue (e.g., Primary Prostate Epithelial Cells, Normal, Human, ATCC PCS-440-010), supplement Prostate Epithelial Cell Basal Medium with the Prostate Epithelial Cell Growth Kit (ATCC PCS-440-040).
- 2. Optional media supplements:
 - 1. Gentamicin-Amphotericin B Solution (ATCC PCS-999-025)
 - 2. Penicillin-Streptomycin-Amphotericin B Solution (ATCC PCS-999-002)
 - 3. Phenol Red (ATCC PCS-999-001)

Volume: 485 ml



Directions for Use

- 1. Obtain one growth kit from the freezer; make sure that the caps of all components are tight.
- Thaw the components of the growth kit just prior to adding them to the basal medium. If the growth kit contains L-glutamine, warm the L-glutamine component in a 37°C water bath and shake to dissolve any precipitates prior to adding to the basal medium.
- 3. Obtain one bottle of Prostate Cell Basal Medium (485 mL) from cold storage.
- Decontaminate the external surfaces of all growth kit component vials and the basal medium bottle by spraying them with 70% ethanol.
- 5. Using aseptic technique and working in a laminar flow hood or biosafety cabinet, transfer the indicated volume of each growth kit component, as indicated in Table 1, to the bottle of basal medium using a separate sterile pipette for each transfer.

Table 1. Prostate Epithelial Cell Growth Kit Components

Component	Volume	Final
		Concentration
L-Glutamine	15 mL	6 mM
Extract P	2.0 mL	0.4%
Epinephrine	0.5 mL	1.0 mM
rh TGF-a	0.5 mL	0.5 ng/mL
Hydrocortisone	0.5 mL	100 ng/mL
hemisuccinate		
rh Insulin	0.5 mL	5 mg/mL
Apo-transferrin	0.5 mL	5 mg/mL

Antimicrobials and phenol red are not required for proliferation but may be added if desired. The recommended volume of each *optional* component to be added to the complete media is summarized in Table 2.

Table 2. Addition of Antimicrobials/Antibiotics and Phenol Red (Optional)

Component	Volume	Final Concentration
Gentamicin-	0.5 mL	Gentamicin: 10 μg/mL
Amphotericin B		Amphotericin B: 0.25
Solution		μg/mL
Penicillin-	0.5 mL	Penicillin: 10 Units/mL
Streptomycin-		Streptomycin: 10 µg/mL
Amphotericin B		Amphotericin B: 25 ng/mL
Solution		
Phenol Red	0.5 mL	33 µM

- Tightly cap the bottle of complete growth medium and swirl the contents gently to assure a homogeneous solution. Do not shake forcefully to avoid foaming. Label and date the bottle.
- Complete growth media should be stored in the dark at 2°C to 8°C (do not freeze). When stored under these conditions, complete media is stable for 30 days.

Quality Control Specifications

Cell testing: Rate of proliferation and morphology.


pH: 7.5 ±0.2



Product Sheet

Prostate Epithelial Cell Basal Medium (ATCC® PCS-440-030™)

Please read this FIRST



ATCC Warranty

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2013. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [09/09]

American Type Culture Collection PO Box 1549 Manassas, VA 20108 USA www.atcc.org

800.638.6597 or 703.365.2700 Fax: 703.365.2750 Email: Tech@atcc.org

Or contact your local distributor